

(FILE 'HOME' ENTERED AT 16:04:39 ON 16 JAN 2007)

10/529569
MFC 11/4

FILE 'REGISTRY' ENTERED AT 16:04:52 ON 16 JAN 2007

L1 STRUCTURE UPLOADED

=> d l1

L1 HAS NO ANSWERS

L1 STR

* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *

Structure attributes must be viewed using STN Express query preparation.

=> s l1

SAMPLE SEARCH INITIATED 16:05:29 FILE 'REGISTRY'

SAMPLE SCREEN SEARCH COMPLETED - 10 TO ITERATE

100.0% PROCESSED 10 ITERATIONS

0 ANSWERS

SEARCH TIME: 00.00.01

FULL FILE PROJECTIONS: ONLINE **COMPLETE**

BATCH **COMPLETE**

PROJECTED ITERATIONS: 11 TO 389

PROJECTED ANSWERS: 0 TO 0

L2 0 SEA SSS SAM L1

=> s l1 full

FULL SEARCH INITIATED 16:05:36 FILE 'REGISTRY'

FULL SCREEN SEARCH COMPLETED - 214 TO ITERATE

100.0% PROCESSED 214 ITERATIONS

0 ANSWERS

SEARCH TIME: 00.00.01

L3 0 SEA SSS FUL L1

=> fil caplus

COST IN U.S. DOLLARS

SINCE FILE

TOTAL

ENTRY

SESSION

FULL ESTIMATED COST

172.55

172.76

FILE 'CAPLUS' ENTERED AT 16:05:52 ON 16 JAN 2007

USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.

PLEASE SEE "HELP USAGETERMS" FOR DETAILS.

COPYRIGHT (C) 2007 AMERICAN CHEMICAL SOCIETY (ACS)

Copyright of the articles to which records in this database refer is held by the publishers listed in the PUBLISHER (PB) field (available for records published or updated in Chemical Abstracts after December 26, 1996), unless otherwise indicated in the original publications. The CA Lexicon is the copyrighted intellectual property of the American Chemical Society and is provided to assist you in searching databases on STN. Any dissemination, distribution, copying, or storing of this information, without the prior written consent of CAS, is strictly prohibited.

FILE COVERS 1907 - 16 Jan 2007 VOL 146 ISS 4

FILE LAST UPDATED: 15 Jan 2007 (20070115/ED)

Effective October 17, 2005, revised CAS Information Use Policies apply. They are available for your review at:

http://www.cas.org/infopolicy.html

10/529569

FIG 12/4

=> s doppelt, pascal/au
L4 33 DOPPELT, PASCAL/AU

=> s 14 and "fluorine-free"
105016 "FLUORINE"
1288898 "FREE"
456 "FLUORINE-FREE"
("FLUORINE" (W) "FREE")
L5 1 L4 AND "FLUORINE-FREE"

=> d bib abs

L5 ANSWER 1 OF 1 CAPLUS COPYRIGHT 2007 ACS on STN
AN 2004:271188 CAPLUS
DN 140:295114
TI New fluorine-free metal complexes for gas-phase
chemical deposition of metals
IN Doppelt, Pascal
PA Centre National de la Recherche Scientifique CNRS, Fr.
SO Fr. Demande, 21 pp.
CODEN: FRXXBL
DT Patent
LA French
FAN.CNT 1



	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	FR 2845088	A1	20040402	FR 2002-12059	20020930
	FR 2845088	B1	20041203		
	CA 2500386	A1	20040408	CA 2003-2500386	20030925
	WO 2004029061	A1	20040408	WO 2003-FR2820	20030925
	W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW				
	RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
	AU 2003276374	A1	20040419	AU 2003-276374	20030925
	EP 1551851	A1	20050713	EP 2003-798226	20030925
	EP 1551851	B1	20060802		
	R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, HU, SK				
	JP 2006501280	T	20060112	JP 2004-539140	20030925
	AT 334990	T	20060815	AT 2003-798226	20030925
	US 2006121709	A1	20060608	US 2005-529569	20050929
PRAI	FR 2002-12059	A	20020930		
	WO 2003-FR2820	W	20030925		
OS	MARPAT 140:295114				
GI					

M+1 (R'-CO-C(R)-CO-R')-1 .L I

AB The invention has as an aim of new complexes of copper(I) or silver(I) and their use for chemical plating in gas phase of copper or silver metals practically free from impurities, complexes of structure I, in which M is

10/529569

PN6 3/4





10/529569

Cu or Ag; R' and R'', identical or different, represent a group chosen from among a C1-8 alkyl, a -OR''' group, in which R''' is C1-8 alkyl; R is a group chosen from among OR''', in which R''' is C1-8 alkyl, a nitro group NO2, an aldehyde function -CHO, an ester function -COOR'', in which R'' is C1-8 alkyl, and L is a ligand of stabilization.

RE.CNT 1 THERE ARE 1 CITED REFERENCES AVAILABLE FOR THIS RECORD
ALL CITATIONS AVAILABLE IN THE RE FORMAT

PNG 4/4



chain nodes :

10 11 13 14 15 17 21

ring nodes :

1 2 3 4 5 6

chain bonds :

3-10 4-21 5-11 13-14 14-15 15-17

ring bonds :

1-2 1-6 2-3 3-4 4-5 5-6

exact/norm bonds :

1-2 1-6 2-3 3-4 3-10 4-5 4-21 5-6 5-11 13-14 14-15 15-17

G1:Ag,Cu

G2:CH3,Et,n-Pr,i-Pr,n-Bu,i-Bu,s-Bu,t-Bu,MeO,EtO,n-PrO,i-PrO,n-BuO,i-BuO,s-BuO,t-BuO,PhO

G3:MeO,EtO,n-PrO,i-PrO,n-BuO,i-BuO,s-BuO,t-BuO,PhO,CHO,NO2,[*1]

G4:CH3,Et,n-Pr,i-Pr,n-Bu,i-Bu,s-Bu,t-Bu,Ph

Match level :

1:Atom 2:Atom 3:Atom 4:Atom 5:Atom 6:Atom 10:CLASS11:CLASS13:CLASS14:CLASS15:CLASS17:CLASS21:CLASS

L1 STRUCTURE UPLOADED

=> dis l1

L1 HAS NO ANSWERS

L1 STR

*** STRUCTURE DIAGRAM IS NOT AVAILABLE ***

Structure attributes must be viewed using STN Express query preparation.

=> s l1 sam

SAMPLE SEARCH INITIATED 17:52:56 FILE 'REGISTRY'

SAMPLE SCREEN SEARCH COMPLETED - 36 TO ITERATE

100.0% PROCESSED 36 ITERATIONS

2 ANSWERS

SEARCH TIME: 00.00.01

FULL FILE PROJECTIONS: ONLINE **COMPLETE**

BATCH **COMPLETE**

PROJECTED ITERATIONS: 360 TO 1080

PROJECTED ANSWERS: 2 TO 124

L2 2 SEA SSS SAM L1

=> dis bib abs

'BIB' IS NOT A VALID FORMAT FOR FILE 'REGISTRY'

'ABS' IS NOT A VALID FORMAT FOR FILE 'REGISTRY'

The following are valid formats:

Substance information can be displayed by requesting individual fields or predefined formats. The predefined substance formats are: (RN = CAS Registry Number)

REG - RN
SAM - Index Name, MF, and structure - no RN
FIDE - All substance data, except sequence data
IDE - FIDE, but only 50 names
SQIDE - IDE, plus sequence data
SQIDE3 - Same as SQIDE, but 3-letter amino acid codes are used
SQD - Protein sequence data, includes RN
SQD3 - Same as SQD, but 3-letter amino acid codes are used
SQN - Protein sequence name information, includes RN

CALC - Table of calculated properties
EPROP - Table of experimental properties
PROP - EPROP and CALC

Any CA File format may be combined with any substance format to obtain CA references citing the substance. The substance formats must be cited first. The CA File predefined formats are:

ABS -- Abstract
APPS -- Application and Priority Information
BIB -- CA Accession Number, plus Bibliographic Data
CAN -- CA Accession Number
CBIB -- CA Accession Number, plus Bibliographic Data (compressed)

IND -- Index Data
IPC -- International Patent Classification
PATS -- PI, SO
STD -- BIB, IPC, and NCL

IABS -- ABS, indented, with text labels
IBIB -- BIB, indented, with text labels
ISTD -- STD format, indented

OBIB ----- AN, plus Bibliographic Data (original)
OIBIB ----- OBIB, indented with text labels

SBIB ----- BIB, no citations
SIBIB ----- IBIB, no citations

The ALL format gives FIDE BIB ABS IND RE, plus sequence data when it is available.


The MAX format is the same as ALL.

The IALL format is the same as ALL with BIB ABS and IND indented, with text labels.

For additional information, please consult the following help messages:

HELP DFIELDS -- To see a complete list of individual display fields.
HELP FORMATS -- To see detailed descriptions of the predefined formats.
ENTER DISPLAY FORMAT (IDE):ide

L2 ANSWER 1 OF 2 REGISTRY COPYRIGHT 2007 ACS on STN
RN 170659-62-8 REGISTRY
ED Entered STN: 28 Nov 1995
CN Copper(1+), (3-nitro-2,4-pentanedionato-O2,O4) (1-pyrrolidineethanamine-N α ,N1)-, (SP-4-3)- (9CI) (CA INDEX NAME)
MF C11 H20 Cu N3 O4
CI CCS, COM
SR CA



=> dis his

(FILE 'HOME' ENTERED AT 17:51:45 ON 16 JAN 2007)

FILE 'REGISTRY' ENTERED AT 17:51:56 ON 16 JAN 2007

L1 STRUCTURE UPLOADED

L2 2 S L1 SAM

=> dis 12

L2 ANSWER 1 OF 2 REGISTRY COPYRIGHT 2007 ACS on STN

RN 170659-62-8 REGISTRY


ED Entered STN: 28 Nov 1995

CN Copper(1+), (3-nitro-2,4-pentanedionato-O2,O4) (1-pyrrolidineethanamine-N α ,N1)-, (SP-4-3)- (9CI) (CA INDEX NAME)

MF C11 H20 Cu N3 O4

CI CCS, COM

SR CA



=> dis 12 1-2

L2 ANSWER 1 OF 2 REGISTRY COPYRIGHT 2007 ACS on STN

RN 170659-62-8 REGISTRY


ED Entered STN: 28 Nov 1995

CN Copper(1+), (3-nitro-2,4-pentanedionato-O2,O4) (1-pyrrolidineethanamine-N α ,N1)-, (SP-4-3)- (9CI) (CA INDEX NAME)


MF C11 H20 Cu N3 O4

CI CCS, COM

SR CA



L2 ANSWER 2 OF 2 REGISTRY COPYRIGHT 2007 ACS on STN
 RN 65588-32-1 REGISTRY
 ED Entered STN: 16 Nov 1984
 CN Copper, [2-(1-iminoethyl)-4-nitrophenolato-N2,O1](3-nitro-2,4-pentanedionato-O2,O4)- (9CI) (CA INDEX NAME)
 OTHER CA INDEX NAMES:
 CN 2,4-Pentanedione, 3-nitro-, copper complex
 CN Phenol, 2-(1-iminoethyl)-4-nitro-, copper complex
 MF C13 H13 Cu N3 O7
 CI CCS
 LC STN Files: CA, CAPLUS



1 REFERENCES IN FILE CA (1907 TO DATE)
 1 REFERENCES IN FILE CAPLUS (1907 TO DATE)

=> s l1 full
 FULL SEARCH INITIATED 17:54:50 FILE 'REGISTRY'
 FULL SCREEN SEARCH COMPLETED - 675 TO ITERATE


100.0% PROCESSED 675 ITERATIONS
 SEARCH TIME: 00.00.01

49 ANSWERS

L3 49 SEA SSS FUL L1

=> d l3 1-49

L3 ANSWER 1 OF 49 REGISTRY COPYRIGHT 2007 ACS on STN
 RN 675200-72-3 REGISTRY
 ED Entered STN: 14 Apr 2004
 CN Copper, [(η2-1,2-ethynediyl)bis(trimethylsilane)](3-methoxy-2,4-pentanedionato-κO,κO')- (9CI) (CA INDEX NAME)
 MF C14 H27 Cu O3 Si2
 CI CCS
 SR CA
 LC STN Files: CA, CAPLUS, USPATFULL




PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT

1 REFERENCES IN FILE CA (1907 TO DATE)

1 REFERENCES IN FILE CAPLUS (1907 TO DATE)

L3 ANSWER 2 OF 49 REGISTRY COPYRIGHT 2007 ACS on STN
 RN 675200-71-2 REGISTRY
 ED Entered STN: 14 Apr 2004
 CN Copper, [(η²-1,2-ethynediyl)bis(trimethylsilane)](4-nitro-3,5-heptanedionato-κO,κO')- (9CI) (CA INDEX NAME)
 MF C15 H28 Cu N O4 Si2
 CI CCS
 SR CA
 LC STN Files: CA, CAPLUS, USPATFULL




PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT

1 REFERENCES IN FILE CA (1907 TO DATE)

1 REFERENCES IN FILE CAPLUS (1907 TO DATE)


L3 ANSWER 3 OF 49 REGISTRY COPYRIGHT 2007 ACS on STN
 RN 675200-70-1 REGISTRY
 ED Entered STN: 14 Apr 2004
 CN Copper, [(1,2-η)-1,5-cyclooctadiene](3-nitro-2,4-pentanedionato-κO,κO')- (9CI) (CA INDEX NAME)
 MF C13 H18 Cu N O4
 CI CCS
 SR CA
 LC STN Files: CA, CAPLUS, USPATFULL



PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT

1 REFERENCES IN FILE CA (1907 TO DATE)
1 REFERENCES IN FILE CAPLUS (1907 TO DATE)


L3 ANSWER 4 OF 49 REGISTRY COPYRIGHT 2007 ACS on STN
RN 675200-69-8 REGISTRY
ED Entered STN: 14 Apr 2004
CN Copper, [(η²-1,2-ethynediyl)bis(trimethylsilane)](3-nitro-2,4-pentanedionato-κO,κO')- (9CI) (CA INDEX NAME)
MF C13 H24 Cu N O4 Si2
CI CCS
SR CA
LC STN Files: CA, CAPLUS, USPATFULL



PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT

1 REFERENCES IN FILE CA (1907 TO DATE)
1 REFERENCES IN FILE CAPLUS (1907 TO DATE)


L3 ANSWER 5 OF 49 REGISTRY COPYRIGHT 2007 ACS on STN
RN 675200-68-7 REGISTRY
ED Entered STN: 14 Apr 2004
CN Copper, bis(4-nitro-3,5-heptanedionato-κO,κO')- (9CI) (CA INDEX NAME)
MF C14 H20 Cu N2 O8
CI CCS
SR CA
LC STN Files: CA, CAPLUS, USPATFULL



PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT


1 REFERENCES IN FILE CA (1907 TO DATE)
1 REFERENCES IN FILE CAPLUS (1907 TO DATE)

L3 ANSWER 6 OF 49 REGISTRY COPYRIGHT 2007 ACS on STN
RN 461031-90-3 REGISTRY
ED Entered STN: 14 Oct 2002
CN Copper, bis[methyl 2-(acetyl-κO)-3-(oxo-κO)butanoato]- (9CI)
(CA INDEX NAME)
MF C14 H18 Cu O8
CI CCS
SR CA
LC STN Files: CA, CAPLUS




1 REFERENCES IN FILE CA (1907 TO DATE)
1 REFERENCES IN FILE CAPLUS (1907 TO DATE)

L3 ANSWER 7 OF 49 REGISTRY COPYRIGHT 2007 ACS on STN
RN 342805-83-8 REGISTRY
ED Entered STN: 21 Jun 2001
CN Copper, bis(trimethyl methanetricarboxylato-κO''',κO''')-,
(SP-4-1)- (9CI) (CA INDEX NAME)
MF C14 H18 Cu O12
CI CCS
SR CA
LC STN Files: CA, CAPLUS, CASREACT




1 REFERENCES IN FILE CA (1907 TO DATE)
1 REFERENCES IN FILE CAPLUS (1907 TO DATE)

L3 ANSWER 8 OF 49 REGISTRY COPYRIGHT 2007 ACS on STN
RN 342805-82-7 REGISTRY
ED Entered STN: 21 Jun 2001
CN Copper, bis(tetrahydrofuran)bis(trimethyl methanetricarboxylato- $\kappa O'''', \kappa O'''')$ -, (OC-6-11)- (9CI) (CA INDEX NAME)
MF C22 H34 Cu O14
CI CCS
SR CA
LC STN Files: CA, CAPLUS, CASREACT




1 REFERENCES IN FILE CA (1907 TO DATE)
1 REFERENCES IN FILE CAPLUS (1907 TO DATE)

L3 ANSWER 9 OF 49 REGISTRY COPYRIGHT 2007 ACS on STN
RN 342805-80-5 REGISTRY
ED Entered STN: 21 Jun 2001
CN Copper, methoxy(trimethyl methanetricarboxylato- $\kappa O'''', \kappa O'''')$ -, (9CI) (CA INDEX NAME)
MF C8 H12 Cu O7
CI CCS
SR CA
LC STN Files: CA, CAPLUS, CASREACT




1 REFERENCES IN FILE CA (1907 TO DATE)
1 REFERENCES IN FILE CAPLUS (1907 TO DATE).

L3 ANSWER 10 OF 49 REGISTRY COPYRIGHT 2007 ACS on STN
RN 306770-10-5 REGISTRY
ED Entered STN: 05 Dec 2000
CN Copper, [2-(acetyl-κO)-3-(oxo-κO)butanalato] (3-acetyl-2,4-pentanedionato-κO,κO')- (9CI) (CA INDEX NAME)
MF C13 H16 Cu O6
CI CCS
SR Chemical Library
Supplier: Interbioscreen Ltd.
LC STN Files: CHEMCATS




L3 ANSWER 11 OF 49 REGISTRY COPYRIGHT 2007 ACS on STN
RN 207454-72-6 REGISTRY
ED Entered STN: 21 Jun 1998
CN Copper, [α-(methoxy-κO)-α-(2-pyridinyl-κN)-2-pyridinemethanol-κN1] (3-nitro-2,4-pentanedionato-κO,κO') (perchlorato-κO)-, (OC-6-34)- (9CI) (CA INDEX NAME)
MF C17 H18 Cl Cu N3 O10
CI CCS
SR CA
LC STN Files: CA, CAPLUS



1 REFERENCES IN FILE CA (1907 TO DATE)
1 REFERENCES IN FILE CAPLUS (1907 TO DATE)

L3 ANSWER 12 OF 49 REGISTRY COPYRIGHT 2007 ACS on STN
RN 207454-65-7 REGISTRY
ED Entered STN: 21 Jun 1998
CN Copper, aqua(3-nitro-2,4-pentanedionato-κO,κO')(perchlorato-κO)[N-(2-pyridinyl-κN)-2-pyridinamine-κN1]-, (OC-6-34)-
(9CI) (CA INDEX NAME)
MF C15 H17 Cl Cu N4 O9
CI CCS
SR CA
LC STN Files: CA, CAPLUS




1 REFERENCES IN FILE CA (1907 TO DATE)
1 REFERENCES IN FILE CAPLUS (1907 TO DATE)

L3 ANSWER 13 OF 49 REGISTRY COPYRIGHT 2007 ACS on STN
RN 170659-73-1 REGISTRY
ED Entered STN: 28 Nov 1995

CN Copper(1+), (2,9-dimethyl-1,10-phenanthroline-N1,N10) (3-nitro-2,4-pentanedionato-O2,O4)-, perchlorate (9CI) (CA INDEX NAME)
 MF C19 H18 Cu N3 O4 . Cl O4
 SR CA
 LC STN Files: CA, CAPLUS


CM 1

CRN 170659-72-0
 CMF C19 H18 Cu N3 O4
 CCI CCS




CM 2

CRN 14797-73-0
 CMF Cl O4



1 REFERENCES IN FILE CA (1907 TO DATE)
 1 REFERENCES IN FILE CAPLUS (1907 TO DATE)

L3 ANSWER 14 OF 49 REGISTRY COPYRIGHT 2007 ACS on STN
 RN 170659-72-0 REGISTRY
 ED Entered STN: 28 Nov 1995
 CN Copper(1+), (2,9-dimethyl-1,10-phenanthroline-N1,N10) (3-nitro-2,4-pentanedionato-O2,O4)- (9CI) (CA INDEX NAME)
 MF C19 H18 Cu N3 O4
 CI CCS, COM
 SR CA




L3 ANSWER 15 OF 49 REGISTRY COPYRIGHT 2007 ACS on STN
 RN 170659-71-9 REGISTRY
 ED Entered STN: 28 Nov 1995
 CN Copper(1+), (3-nitro-2,4-pentanedionato-O2,O4) (1,10-phenanthroline-N1,N10)-
 , (SP-4-2)-, perchlorate (9CI) (CA INDEX NAME)
 MF C17 H14 Cu N3 O4 . Cl O4
 SR CA
 LC STN Files: CA, CAPLUS

CM 1

CRN 170659-70-8

CMF C17 H14 Cu N3 O4


CCI CCS



CM 2


CRN 14797-73-0

CMF Cl O4



1 REFERENCES IN FILE CA (1907 TO DATE)
1 REFERENCES IN FILE CAPLUS (1907 TO DATE)


L3 ANSWER 16 OF 49 REGISTRY COPYRIGHT 2007 ACS on STN
RN ~~170659-70-8~~ REGISTRY
ED Entered STN: 28 Nov 1995
CN Copper(1+), (3-nitro-2,4-pentanedionato-O2,O4) (1,10-phenanthroline-N1,N10)-
, (SP-4-2)- (9CI) (CA INDEX NAME)
MF C17 H14 Cu N3 O4
CI CCS, COM
SR CA



L3 ANSWER 17 OF 49 REGISTRY COPYRIGHT 2007 ACS on STN
RN 170659-69-5 REGISTRY
ED Entered STN: 28 Nov 1995
CN Copper(1+), (2,2'-bipyridine-N,N') (3-nitro-2,4-pentanedionato-O2,O4)-,
(SP-4-3)-, perchlorate (9CI) (CA INDEX NAME)
MF C15 H14 Cu N3 O4 . Cl O4
SR CA
LC STN Files: CA, CAPLUS

CM 1


CRN 170659-68-4
CMF C15 H14 Cu N3 O4
CCI CCS



CM 2

CRN 14797-73-0


CMF Cl 04



1 REFERENCES IN FILE CA (1907 TO DATE)

1 REFERENCES IN FILE CAPLUS (1907 TO DATE)

L3 ANSWER 18 OF 49 REGISTRY COPYRIGHT 2007 ACS on STN
 RN 170659-68-4 REGISTRY
 ED Entered STN: 28 Nov 1995
 CN Copper(1+), (2,2'-bipyridine-N,N') (3-nitro-2,4-pentanedionato-02,04)-,
 (SP-4-3)- (9CI) (CA INDEX NAME)
 MF C15 H14 Cu N3 O4
 CI CCS, COM
 SR CA




L3 ANSWER 19 OF 49 REGISTRY COPYRIGHT 2007 ACS on STN

RN 170659-67-3 REGISTRY
 ED Entered STN: 28 Nov 1995
 CN Copper(1+), (3-nitro-2,4-pentanedionato-O2,O4) (2-pyridinemethanamine-N1,N2)-, (SP-4-3)-, perchlorate (9CI) (CA INDEX NAME)
 MF C11 H14 Cu N3 O4 . Cl O4
 SR CA
 LC STN Files: CA, CAPLUS


CM 1

CRN 170659-66-2
 CMF C11 H14 Cu N3 O4
 CCI CCS




CM 2

CRN 14797-73-0
 CMF Cl O4



1 REFERENCES IN FILE CA (1907 TO DATE)
 1 REFERENCES IN FILE CAPLUS (1907 TO DATE)


L3 ANSWER 20 OF 49 REGISTRY COPYRIGHT 2007 ACS on STN
 RN 170659-66-2 REGISTRY
 ED Entered STN: 28 Nov 1995
 CN Copper(1+), (3-nitro-2,4-pentanedionato-O2,O4) (2-pyridinemethanamine-N1,N2)-, (SP-4-3)- (9CI) (CA INDEX NAME)
 MF C11 H14 Cu N3 O4
 CI CCS, COM
 SR CA



L3 ANSWER 21 OF 49 REGISTRY COPYRIGHT 2007 ACS on STN
 RN 170659-65-1 REGISTRY
 ED Entered STN: 28 Nov 1995
 CN Copper(1+), (1-ethyl-2-pyrrolidinemethanamine-N α ,N1) (3-nitro-2,4-pentanedionato-O2,O4)-, (SP-4-3)-, perchlorate (9CI) (CA INDEX NAME)
 MF C12 H22 Cu N3 O4 . Cl O4
 SR CA
 LC STN Files: CA, CAPLUS


CM 1

CRN 170659-64-0
 CMF C12 H22 Cu N3 O4
 CCI CCS




CM 2

CRN 14797-73-0
 CMF Cl O4



1 REFERENCES IN FILE CA (1907 TO DATE)
1 REFERENCES IN FILE CAPLUS (1907 TO DATE)


L3 **ANSWER 22 OF 49** REGISTRY COPYRIGHT 2007 ACS on STN
RN 170659-64-0 REGISTRY
ED Entered STN: 28 Nov 1995
CN Copper(1+), (1-ethyl-2-pyrrolidinemethanamine-N α ,N1) (3-nitro-2,4-pentanedionato-02,04)-, (SP-4-3)- (9CI) (CA INDEX NAME)
MF C12 H22 Cu N3 O4
CI CCS, COM
SR CA



L3 **ANSWER 23 OF 49** REGISTRY COPYRIGHT 2007 ACS on STN
RN 170659-63-9 REGISTRY
ED Entered STN: 28 Nov 1995
CN Copper(1+), (3-nitro-2,4-pentanedionato-02,04) (1-pyrrolidineethanamine-N α ,N1)-, (SP-4-3)-, perchlorate (9CI) (CA INDEX NAME)
MF C11 H20 Cu N3 O4 . Cl O4
SR CA
LC STN Files: CA, CAPLUS

CM 1


CRN 170659-62-8
CMF C11 H20 Cu N3 O4
CCI CCS



CM 2

CRN 14797-73-0


CMF Cl O4



1 REFERENCES IN FILE CA (1907 TO DATE)

1 REFERENCES IN FILE CAPLUS (1907 TO DATE)


L3 ANSWER 24 OF 49 REGISTRY COPYRIGHT 2007 ACS on STN
 RN 170659-62-8 REGISTRY
 ED Entered STN: 28 Nov 1995
 CN Copper(1+), (3-nitro-2,4-pentanedionato-O2,O4) (1-pyrrolidineethanamine-Na,N1)-, (SP-4-3)- (9CI) (CA INDEX NAME)
 MF C11 H20 Cu N3 O4
 CI CCS, COM
 SR CA



L3 ANSWER 25 OF 49 REGISTRY COPYRIGHT 2007 ACS on STN
 RN 152767-34-5 REGISTRY
 ED Entered STN: 04 Feb 1994
 CN Copper(1+), (N,N-dimethyl-1,2-ethanediamine-κN,κN') (3-nitro-2,4-pentanedionato-κO,κO')-, (SP-4-3)-, nitrate (9CI) (CA INDEX NAME)
 OTHER CA INDEX NAMES:
 CN Copper(1+), (N,N-dimethyl-1,2-ethanediamine-N,N') (3-nitro-2,4-pentanedionato-O2,O4)-, (SP-4-3)-, nitrate
 MF C9 H18 Cu N3 O4 . N O3
 SR CA
 LC STN Files: CA, CAPLUS


CM 1

CRN 152748-67-9
CMF C9 H18 Cu N3 O4
CCI CCS



CM 2

CRN 14797-55-8
CMF N O3




1 REFERENCES IN FILE CA (1907 TO DATE)
1 REFERENCES IN FILE CAPLUS (1907 TO DATE)

L3 ANSWER 26 OF 49 REGISTRY COPYRIGHT 2007 ACS on STN
RN 152748-68-0 REGISTRY
ED Entered STN: 04 Feb 1994
CN Copper(1+), (N,N-dimethyl-1,2-ethanediamine-N,N') (3-nitro-2,4-pentanedionato-O2,O4)-, (SP-4-3)-, perchlorate (9CI) (CA INDEX NAME)
MF C9 H18 Cu N3 O4 . Cl O4
SR CA
LC STN Files: CA, CAPLUS

CM 1


CRN 152748-67-9
CMF C9 H18 Cu N3 O4
CCI CCS



CM 2

CRN 14797-73-0


CMF Cl 04




2 REFERENCES IN FILE CA (1907 TO DATE)

2 REFERENCES IN FILE CAPLUS (1907 TO DATE)

L3 ANSWER 27 OF 49 REGISTRY COPYRIGHT 2007 ACS on STN
 RN 152748-67-9 REGISTRY
 ED Entered STN: 04 Feb 1994
 CN Copper(1+), (N,N-dimethyl-1,2-ethanediamine-N,N')(3-nitro-2,4-pentanedionato-O2,O4)-, (SP-4-3)- (9CI) (CA INDEX NAME)
 MF C9 H18 Cu N3 O4
 CI CCS, COM
 SR CA



L3 ANSWER 28 OF 49 REGISTRY COPYRIGHT 2007 ACS on STN
 RN 148379-28-6 REGISTRY
 ED Entered STN: 30 Jun 1993
 CN Copper(1+), (N'-ethyl-N,N-dimethyl-1,2-ethanediamine-N,N')(3-nitro-2,4-pentanedionato-O2,O4)-, chloride, (SP-4-3)- (9CI) (CA INDEX NAME)
 MF C11 H22 Cu N3 O4 . Cl
 CI CCS
 SR CA
 LC STN Files: CA, CAPLUS
 CRN (148379-25-3)




● Cl⁻

1 REFERENCES IN FILE CA (1907 TO DATE)
1 REFERENCES IN FILE CAPLUS (1907 TO DATE)

L3 **ANSWER 29 OF 49** REGISTRY COPYRIGHT 2007 ACS on STN
RN 148379-27-5 REGISTRY
ED Entered STN: 30 Jun 1993
CN Copper(1+), (N'-ethyl-N,N-dimethyl-1,2-ethanediamine-N,N') (3-nitro-2,4-pentanedionato-O2,O4)-, (SP-4-3)-, nitrate (9CI) (CA INDEX NAME)
MF C11 H22 Cu N3 O4 . N O3
SR CA
LC STN Files: CA, CAPLUS


CM 1

CRN 148379-25-3
CMF C11 H22 Cu N3 O4
CCI CCS



CM 2

CRN 14797-55-8
CMF N O3




1 REFERENCES IN FILE CA (1907 TO DATE)
1 REFERENCES IN FILE CAPLUS (1907 TO DATE)

L3 ANSWER 30 OF 49 REGISTRY COPYRIGHT 2007 ACS on STN
RN 148379-26-4 REGISTRY
ED Entered STN: 30 Jun 1993
CN Copper(1+), (N'-ethyl-N,N-dimethyl-1,2-ethanediamine-N,N')(3-nitro-2,4-pentanedionato-O2,O4)-, (SP-4-3)-, perchlorate (9CI) (CA INDEX NAME)
MF C11 H22 Cu N3 O4 . Cl O4
SR CA
LC STN Files: CA, CAPLUS


CM 1

CRN 148379-25-3
CMF C11 H22 Cu N3 O4
CCI CCS




CM 2

CRN 14797-73-0
CMF Cl O4




1 REFERENCES IN FILE CA (1907 TO DATE)
1 REFERENCES IN FILE CAPLUS (1907 TO DATE)

L3 ANSWER 31 OF 49 REGISTRY COPYRIGHT 2007 ACS on STN
RN 148379-25-3 REGISTRY
ED Entered STN: 30 Jun 1993
CN Copper(1+), (N'-ethyl-N,N-dimethyl-1,2-ethanediamine-N,N')(3-nitro-2,4-pentanedionato-O2,O4)-, (SP-4-3)- (9CI) (CA INDEX NAME)
MF C11 H22 Cu N3 O4
CI CCS, COM
SR CA




L3 **ANSWER 32 OF 49** REGISTRY COPYRIGHT 2007 ACS on STN
 RN 138072-61-4 REGISTRY
 ED Entered STN: 27 Dec 1991
 CN Copper, bis(1,4-butanediamine-N)bis(3-nitro-2,4-pentanedionato-O2,O4)-
 (9CI) (CA INDEX NAME)
 MF C18 H36 Cu N6 O8
 CI CCS
 SR CA
 LC STN Files: CA, CAPLUS




3 REFERENCES IN FILE CA (1907 TO DATE)
 3 REFERENCES IN FILE CAPLUS (1907 TO DATE)

L3 **ANSWER 33 OF 49** REGISTRY COPYRIGHT 2007 ACS on STN
 RN 138055-62-6 REGISTRY
 ED Entered STN: 27 Dec 1991
 CN Copper, bis(3-nitro-2,4-pentanedionato-O2,O4)bis(1,3-propanediamine-N)-
 (9CI) (CA INDEX NAME)
 MF C16 H32 Cu N6 O8
 CI CCS
 SR CA
 LC STN Files: CA, CAPLUS




4 REFERENCES IN FILE CA (1907 TO DATE)
4 REFERENCES IN FILE CAPLUS (1907 TO DATE)

L3 **ANSWER 34 OF 49** REGISTRY COPYRIGHT 2007 ACS on STN
RN 128303-88-8 REGISTRY
ED Entered STN: 20 Jul 1990
CN Acetoacetaldehyde, 2-acetyl-, Cu deriv. (6CI) (CA INDEX NAME)
MF C12 H14 Cu O6
CI CCS
SR CAOLD
LC STN Files: CAOLD



1 REFERENCES IN FILE CAOLD (PRIOR TO 1967)

L3 **ANSWER 35 OF 49** REGISTRY COPYRIGHT 2007 ACS on STN
RN 118168-01-7 REGISTRY
ED Entered STN: 23 Dec 1988
CN Cuprate(1-), bis(3-nitro-2,4-pentanedionato-O2,O4)-, (T-4)- (9CI) (CA INDEX NAME)
OTHER CA INDEX NAMES:
CN 2,4-Pentanedione, 3-nitro-, copper complex
MF C10 H12 Cu N2 O8
CI CCS
SR CA
LC STN Files: CA, CAPLUS



1 REFERENCES IN FILE CA (1907 TO DATE)
1 REFERENCES IN FILE CAPLUS (1907 TO DATE)

L3 **ANSWER 36 OF 49** REGISTRY COPYRIGHT 2007 ACS on STN
RN 110873-86-4 REGISTRY
ED Entered STN: 24 Oct 1987
CN Copper, bis(1,2-ethanediamine-N)bis(3-nitro-2,4-pentanedionato-O2,O4)- (9CI) (CA INDEX NAME)
OTHER CA INDEX NAMES:

CN 2,4-Pentanedione, 3-nitro-, copper complex

OTHER NAMES:


CN Bis(ethylenediamine)bis(3-nitro-2,4-pentanedionato)copper

MF C14 H28 Cu N6 O8

CI CCS

SR CA

LC STN Files: CA, CAPLUS



4 REFERENCES IN FILE CA (1907 TO DATE)

4 REFERENCES IN FILE CAPLUS (1907 TO DATE)

L3 **ANSWER 37 OF 49** REGISTRY COPYRIGHT 2007 ACS on STN

RN 76401-08-6 REGISTRY

ED Entered STN: 16 Nov 1984

CN Copper, bis(3-phenoxy-2,4-pentanedionato-O2,O4)- (9CI) (CA INDEX NAME)


OTHER CA INDEX NAMES:

CN 2,4-Pentanedione, 3-phenoxy-, copper complex

MF C22 H22 Cu O6

CI CCS

LC STN Files: CA, CAPLUS



1 REFERENCES IN FILE CA (1907 TO DATE)

1 REFERENCES IN FILE CAPLUS (1907 TO DATE)

L3 **ANSWER 38 OF 49** REGISTRY COPYRIGHT 2007 ACS on STN

RN 65888-28-0 REGISTRY

ED Entered STN: 16 Nov 1984

CN Copper, [2-(iminomethyl)-4,6-dinitrophenolato-N2,O1](3-nitro-2,4-pentanedionato-O2,O4)- (9CI) (CA INDEX NAME)

OTHER CA INDEX NAMES:


CN 2,4-Pentanedione, 3-nitro-, copper complex

CN Phenol, 2-(iminomethyl)-4,6-dinitro-, copper complex

MF C12 H10 Cu N4 O9

CI CCS


LC STN Files: CA, CAPLUS



1 REFERENCES IN FILE CA (1907 TO DATE)

1 REFERENCES IN FILE CAPLUS (1907 TO DATE)


L3 ANSWER 39 OF 49 REGISTRY COPYRIGHT 2007 ACS on STN
 RN 65588-32-1 REGISTRY
 ED Entered STN: 16 Nov 1984
 CN Copper, [2-(1-iminoethyl)-4-nitrophenolato-N2,O1](3-nitro-2,4-pentanedionato-O2,O4)- (9CI) (CA INDEX NAME)
 OTHER CA INDEX NAMES:
 CN 2,4-Pentanedione, 3-nitro-, copper complex
 CN Phenol, 2-(1-iminoethyl)-4-nitro-, copper complex
 MF C13 H13 Cu N3 O7
 CI CCS
 LC STN Files: CA, CAPLUS



1 REFERENCES IN FILE CA (1907 TO DATE)

1 REFERENCES IN FILE CAPLUS (1907 TO DATE)


L3 ANSWER 40 OF 49 REGISTRY COPYRIGHT 2007 ACS on STN
 RN 65588-31-0 REGISTRY
 ED Entered STN: 16 Nov 1984
 CN Copper, [2-(1-iminoethyl)-4,6-dinitrophenolato-N2,O1](3-nitro-2,4-pentanedionato-O2,O4)- (9CI) (CA INDEX NAME)
 OTHER CA INDEX NAMES:
 CN 2,4-Pentanedione, 3-nitro-, copper complex
 CN Phenol, 2-(1-iminoethyl)-4,6-dinitro-, copper complex
 MF C13 H12 Cu N4 O9
 CI CCS
 LC STN Files: CA, CAPLUS



1 REFERENCES IN FILE CA (1907 TO DATE)

1 REFERENCES IN FILE CAPLUS (1907 TO DATE)

L3 **ANSWER 41 OF 49** REGISTRY. COPYRIGHT 2007 ACS on STN
 RN 55140-20-0 REGISTRY
 ED Entered STN: 16 Nov 1984
 CN Copper, (4-methylpyridine)bis(3-nitro-2,4-pentanedionato-O2,O4)- (9CI)
 (CA INDEX NAME)
 OTHER CA INDEX NAMES:
 CN 2,4-Pentanedione, 3-nitro-, copper complex
 MF C16 H19 Cu N3 O8
 CI CCS
 LC STN Files: CA, CAPLUS




1 REFERENCES IN FILE CA (1907 TO DATE)

1 REFERENCES IN FILE CAPLUS (1907 TO DATE)


L3 **ANSWER 42 OF 49** REGISTRY. COPYRIGHT 2007 ACS on STN
 RN 53494-99-8 REGISTRY
 ED Entered STN: 16 Nov 1984
 CN Copper, bis(3-nitro-2,4-pentanedionato-κO,κO')-, (SP-4-1)-
 (9CI) (CA INDEX NAME)
 OTHER CA INDEX NAMES:
 CN 2,4-Pentanedione, 3-nitro-, copper complex
 CN Copper, bis(3-nitro-2,4-pentanedionato-O2,O4)-, (SP-4-1)-
 OTHER NAMES:

CN Bis(3-nitro-2,4-pentanedionato)copper
 MF C10 H12 Cu N2 O8
 CI CCS
 LC STN Files: CA, CAPLUS, GMELIN*
 (*File contains numerically searchable property data)



7 REFERENCES IN FILE CA (1907 TO DATE)
 8 REFERENCES IN FILE CAPLUS (1907 TO DATE)


L3 ANSWER 43 OF 49 REGISTRY COPYRIGHT 2007 ACS on STN
 RN 41101-63-7 REGISTRY
 ED Entered STN: 16 Nov 1984
 CN Copper, bis(3-nitro-2,4-pentanedionato-O2,O4)(quinoline)- (9CI) (CA INDEX NAME)
 OTHER CA INDEX NAMES:
 CN 2,4-Pentanedione, 3-nitro-, copper complex
 MF C19 H19 Cu N3 O8
 CI CCS
 LC STN Files: CA, CAPLUS



1 REFERENCES IN FILE CA (1907 TO DATE)
 1 REFERENCES IN FILE CAPLUS (1907 TO DATE)


L3 ANSWER 44 OF 49 REGISTRY COPYRIGHT 2007 ACS on STN
 RN 36364-01-9 REGISTRY
 ED Entered STN: 16 Nov 1984
 CN Copper, bis(3-nitro-2,4-pentanedionato-O2,O4)bis(pyridine)- (9CI) (CA INDEX NAME)
 OTHER CA INDEX NAMES:
 CN 2,4-Pentanedione, 3-nitro-, copper complex
 MF C20 H22 Cu N4 O8
 CI CCS

LC STN Files: CA, CAPLUS



1 REFERENCES IN FILE CA (1907 TO DATE)
1 REFERENCES IN FILE CAPLUS (1907 TO DATE)


L3 **ANSWER 45 OF 49** REGISTRY COPYRIGHT 2007 ACS on STN
RN 36363-99-2 REGISTRY
ED Entered STN: 16 Nov 1984
CN Copper, bis(3-nitro-2,4-pentanedionato-O2,O4)(pyridine)-, (SP-5-21)- (9CI)
(CA INDEX NAME)
OTHER CA INDEX NAMES:
CN 2,4-Pentanedione, 3-nitro-, copper complex
MF C15 H17 Cu N3 O8
CI CCS
LC STN Files: CA, CAPLUS



1 REFERENCES IN FILE CA (1907 TO DATE)
1 REFERENCES IN FILE CAPLUS (1907 TO DATE)

L3 **ANSWER 46 OF 49** REGISTRY COPYRIGHT 2007 ACS on STN
RN 36363-97-0 REGISTRY
ED Entered STN: 16 Nov 1984
CN Copper, bis(ethyl 2-acetyl-3-oxobutanoato-O2,O3)(pyridine)-, (SP-5-21)-
(9CI) (CA INDEX NAME)
OTHER CA INDEX NAMES:
CN Butanoic acid, 2-acetyl-3-oxo-, ethyl ester, copper complex

MF C21 H27 Cu N O8
 CI CCS
 LC STN Files: CA, CAPLUS



1 REFERENCES IN FILE CA (1907 TO DATE)
 1 REFERENCES IN FILE CAPLUS (1907 TO DATE)

L3 ANSWER 47 OF 49 REGISTRY COPYRIGHT 2007 ACS on STN
 RN 27970-19-0 REGISTRY
 ED Entered STN: 16 Nov 1984
 CN Copper, bis(3-methoxy-2,4-pentanedionato- κ O, κ O')- (9CI) (CA INDEX NAME)

OTHER CA INDEX NAMES:

CN Copper, bis(3-methoxy-2,4-pentanedionato)- (7CI, 8CI)


OTHER NAMES:

CN Bis(3-methoxy-2,4-pentanedionato)copper

MF C12 H18 Cu O6

CI CCS


LC STN Files: CA, CAOLD, CAPLUS



4 REFERENCES IN FILE CA (1907 TO DATE)
 4 REFERENCES IN FILE CAPLUS (1907 TO DATE)
 1 REFERENCES IN FILE CAOLD (PRIOR TO 1967)


L3 ANSWER 48 OF 49 REGISTRY COPYRIGHT 2007 ACS on STN
 RN 15378-83-3 REGISTRY
 ED Entered STN: 16 Nov 1984
 CN Copper, bis[ethyl 2-(acetyl- κ O)-3-(oxo- κ O)butanoato]- (9CI) (CA INDEX NAME)
 OTHER CA INDEX NAMES:
 CN Copper, bis(hydrogen 2-acetylacetoacetato)-, diethyl ester (7CI, 8CI)
 OTHER NAMES:

CN Copper bis(3-ethoxycarbonylacetylacetone)
 MF C16 H22 Cu O8
 CI CCS
 LC STN Files: CA, CAOLD, CAPLUS, TOXCENTER



7 REFERENCES IN FILE CA (1907 TO DATE)
 7 REFERENCES IN FILE CAPLUS (1907 TO DATE)
 4 REFERENCES IN FILE CAOLD (PRIOR TO 1967)

L3 ANSWER 49 OF 49 REGISTRY COPYRIGHT 2007 ACS on STN
 RN 14689-25-9 REGISTRY
 ED Entered STN: 16 Nov 1984
 CN Copper, bis(3-nitro-2,4-pentanedionato-O2,O4)- (9CI) (CA INDEX NAME)
 OTHER CA INDEX NAMES:
 CN 2,4-Pentanedione, 3-nitro-, copper complex
 CN Copper, bis(3-nitro-2,4-pentanedionato)- (7CI, 8CI)
 OTHER NAMES:
 CN Bis(3-nitroacetylacetonato)copper
 CN NSC 338135
 MF C10 H12 Cu N2 O8
 CI CCS
 LC STN Files: CA, CAOLD, CAPLUS, GMELIN*, USPATFULL
 (*File contains numerically searchable property data)



PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT

27 REFERENCES IN FILE CA (1907 TO DATE)
 27 REFERENCES IN FILE CAPLUS (1907 TO DATE)
 6 REFERENCES IN FILE CAOLD (PRIOR TO 1967)

=> fil hcaplus
 COST IN U.S. DOLLARS

SINCE FILE TOTAL
 ENTRY SESSION

FULL ESTIMATED COST

277.70 . 277.91

FILE 'HCAPLUS' ENTERED AT 17:55:47 ON 16 JAN 2007
 USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.
 PLEASE SEE "HELP USAGETERMS" FOR DETAILS.
 COPYRIGHT (C) 2007 AMERICAN CHEMICAL SOCIETY (ACS)

Copyright of the articles to which records in this database refer is held by the publishers listed in the PUBLISHER (PB) field (available for records published or updated in Chemical Abstracts after December 26, 1996), unless otherwise indicated in the original publications. The CA Lexicon is the copyrighted intellectual property of the the American Chemical Society and is provided to assist you in searching databases on STN. Any dissemination, distribution, copying, or storing of this information, without the prior written consent of CAS, is strictly prohibited.

FILE COVERS 1907 - 16 Jan 2007 VOL 146 ISS 4
 FILE LAST UPDATED: 15 Jan 2007 (20070115/ED)

New CAS Information Use Policies, enter HELP USAGETERMS for details.

This file contains CAS Registry Numbers for easy and accurate substance identification.

=> s l3/prep

50 L3
 4345600 PREP/RL
 L4 23 L3/PREP
 (L3 (L) PREP/RL)

=> d l4 1-23

L4 **ANSWER 1 OF 23** HCAPLUS COPYRIGHT 2007 ACS on STN
 AN 2004:271188 HCAPLUS
 DN 140:295114
 TI New fluorine-free metal complexes for gas-phase chemical deposition of metals
 IN Doppelt, Pascal
 PA Centre National de la Recherche Scientifique CNRS, Fr.
 SO Fr. Demande, 21 pp.
 CODEN: FRXXBL
 DT Patent
 LA French
 FAN.CNT 1

Application

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	FR 2845088	A1	20040402	FR 2002-12059	20020930
	FR 2845088	B1	20041203		
	CA 2500386	A1	20040408	CA 2003-2500386	20030925
	WO 2004029061	A1	20040408	WO 2003-FR2820	20030925

W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN,

TI Preparation and reduction of bis-(3-nitro-2,4-pentanedione)-beryllium
 AU Klein, Richard M.; Bailar, John C., Jr.
 CS Univ. of Illinois, Urbana
 SO Inorg. Chem. (1963), 2(6), 1187-90
 DT Journal
 LA Unavailable

L4 ANSWER 23 OF 23 HCAPLUS COPYRIGHT 2007 ACS on STN
 AN 1963:18976 HCAPLUS
 DN 58:18976
 OREF 58:3090h,3091a-b
 TI The reactivity of coordinated acetylacetone
 AU Djordjevic, C.; Lewis, J.; Nyholm, R. S.
 CS Univ. Coll., London
 SO Journal of the Chemical Society (1962) 4778-84
 CODEN: JCSOA9; ISSN: 0368-1769
 DT Journal
 LA Unavailable

=> d 14 1-23 bib abs hitstr

L4 ANSWER 1 OF 23 HCAPLUS COPYRIGHT 2007 ACS on STN
 AN 2004:271188 HCAPLUS
 DN 140:295114
 TI New fluorine-free metal complexes for gas-phase chemical deposition of metals
 IN Doppelt, Pascal
 PA Centre National de la Recherche Scientifique CNRS, Fr.
 SO Fr. Demande, 21 pp.
 CODEN: FRXXBL
 DT Patent
 LA French
 FAN.CNT 1

just Application

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	FR 2845088	A1	20040402	FR 2002-12059	20020930
	FR 2845088	B1	20041203		
	CA 2500386	A1	20040408	CA 2003-2500386	20030925
	WO 2004029061	A1	20040408	WO 2003-FR2820	20030925
	W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW				
	RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
	AU 2003276374	A1	20040419	AU 2003-276374	20030925
	EP 1551851	A1	20050713	EP 2003-798226	20030925
	EP 1551851	B1	20060802		
	R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, HU, SK				
	JP 2006501280	T	20060112	JP 2004-539140	20030925
	AT 334990	T	20060815	AT 2003-798226	20030925

US 2006121709 A1 20060608 US 2005-529569 20050929
 PRAI FR 2002-12059 A 20020930
 WO 2003-FR2820 W 20030925
 OS MARPAT 140:295114
 GI

$M^{+1}(R'-CO-C(R)-CO-R'')^{-1} \cdot L \cdot I$

AB The invention has as an aim of new complexes of copper(I) or silver(I) and their use for chemical plating in gas phase of copper or silver metals practically free from impurities, complexes of structure I, in which M is Cu or Ag; R' and R'', identical or different, represent a group chosen from among a C1-8 alkyl, a -OR''' group, in which R''' is C1-8 alkyl; R is a group chosen from among OR''', in which R''' is C1-8 alkyl, a nitro group NO₂, an aldehyde function -CHO, an ester function -COOR''', in which R''' is C1-8 alkyl, and L is a ligand of stabilization.

IT 14689-25-9P 675200-68-7P 675200-69-8P
 675200-70-1P 675200-71-2P 675200-72-3P


RL: RCT (Reactant); SPN (Synthetic preparation); PREP

(Preparation); RACT (Reactant or reagent)

(fluorine-free metal complexes for gas-phase chemical deposition of metals)


RN 14689-25-9 HCAPLUS

CN Copper, bis(3-nitro-2,4-pentanedionato-O₂,O₄)-(9CI) (CA INDEX NAME)




RN 675200-68-7 HCAPLUS

CN Copper, bis(4-nitro-3,5-heptanedionato-κO,κO')-(9CI) (CA INDEX NAME)




RN 675200-69-8 HCAPLUS


CN Copper, [(η²-1,2-ethynediyl)bis(trimethylsilane)](3-nitro-2,4-pentanedionato-κO,κO')-(9CI) (CA INDEX NAME)




RN 675200-70-1 HCAPLUS
 CN Copper, [(1,2- η)-1,5-cyclooctadiene] (3-nitro-2,4-pentanedionato- κ O, κ O')- (9CI) (CA INDEX NAME)



RN 675200-71-2 HCAPLUS
 CN Copper, [(η 2-1,2-ethynediyl)bis(trimethylsilane)] (4-nitro-3,5-heptanedionato- κ O, κ O')- (9CI) (CA INDEX NAME)



RN 675200-72-3 HCAPLUS
 CN Copper, [(η 2-1,2-ethynediyl)bis(trimethylsilane)] (3-methoxy-2,4-pentanedionato- κ O, κ O')- (9CI) (CA INDEX NAME)



RE.CNT 1 THERE ARE 1 CITED REFERENCES AVAILABLE FOR THIS RECORD
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 2 OF 23 HCAPLUS COPYRIGHT 2007 ACS on STN

AN 2001:160846 HCAPLUS

DN 135:28213

TI Synthesis and structural characterization of tricarbomethoxymethanate complexes of copper(II) and barium(II) and evaluation of their suitability for MOCVD applications

AU Baxter, David V.; Caulton, Kenneth G.; Chiang, Wen-Chung; Chisholm, Malcolm H.; DiStasi, Vincent F.; Dutremez, Sylvain G.; Martin, James D.; Streib, William E.

CS Department of Physics, Department of Chemistry and Molecular Structure Center, Indiana University, Bloomington, IN, 47405-7102, USA

SO New Journal of Chemistry (2001), 25(3), 400-407
CODEN: NJCHE5; ISSN: 1144-0546.

PB Royal Society of Chemistry

DT Journal

LA English

OS CASREACT 135:28213

AB The reaction between $\text{Cu}(\text{OCH}_3)_2$ and HL [$\text{L} = -\text{C}(\text{CO}_2\text{CH}_3)_3$] in THF gives a mixture of two products, $\text{CuL}(\text{OCH}_3)$ and $\text{CuL}_2(\text{THF})_2$. $\text{CuL}(\text{OCH}_3)$ was characterized by elemental anal., IR spectroscopy and thermal decomposition studies, and its insol. in THF, at room temperature, suggests that it may be polymeric. The solid state structure of $\text{CuL}_2(\text{THF})_2$ was established via a single-crystal x-ray diffraction anal. This complex has a tetragonally distorted trans-disubstituted octahedral structure with L binding like a conventional bidentate β -diketonate ligand; the central CO_2CH_3 group of each L is not coordinated. $\text{CuL}_2(\text{THF})_2$ readily loses the two THF mols. bound to the Cu(II) center upon drying in a stream of dinitrogen to produce THF-free CuL_2 . Alternatively, THF-free CuL_2 can be prepared by the reaction of $\text{CuL}(\text{OCH}_3)$ with HL in refluxing toluene. X-ray crystallog. shows that CuL_2 is a polymeric solid composed of a planar $\text{Cu}(\eta^2\text{-L})_2$ repeat unit in which the central CO_2CH_3 group of each L binds weakly to the Cu(II) center of a neighboring unit along a "stepped" (displaced) stack of CuO_4 units. The reaction between elemental barium and HL in THF requires activation with $\text{NH}_3(\text{g})$. The product is a white solid of empirical formula $\text{BaL}_2 \cdot 0.36\text{THF}$. It was characterized by elemental anal., ^1H NMR, $^{13}\text{C}\{^1\text{H}\}$ NMR, and IR spectroscopies, and thermal decomposition studies. It is insol. in common laboratory solvents but soluble in strong

Lewis

bases such as pyridine, DMSO, and hexamethylphosphoramide (HMPA). The product resulting from the depolymn. of $\text{BaL}_2 \cdot 0.36\text{THF}$ with HMPA was characterized via a single-crystal x-ray diffraction anal. It is a dimer of empirical formula $\text{Ba}_2\text{L}_4(\text{HMPA})_4$. It consists of two $\text{BaL}_2(\text{HMPA})_2$ fragments linked together, in a centrosym. fashion, by two μ_2, η^3 - $[-\text{C}(\text{CO}_2\text{CH}_3)_3]$ ligands; this is made possible using the donor power of the pendant $\text{C}(\text{O})\text{OCH}_3$ group not used in $\text{CuL}_2(\text{THF})_2$. The remaining two $-\text{C}(\text{CO}_2\text{CH}_3)_3$ ligands are also bidentate but are nonbridging, and the four HMPA mols. are bonded through oxygen, resulting in a coordination number of seven for each barium. CuL_2 is volatile and sublimes upon heating the amorphous material obtained from spontaneous desolvation of $\text{CuL}_2(\text{THF})_2$, or polymeric CuL_2 , or $\text{CuL}(\text{OCH}_3)$ under high vacuum. The barium complexes are not volatile. Metallic copper was obtained upon heating the Cu(II) complexes to 1000° in a stream of argon, and formation of CuO results when these complexes are decomposed in the presence of oxygen. Thermal decomposition of $\text{BaL}_2 \cdot 0.36\text{THF}$ and $\text{Ba}_2\text{L}_4(\text{HMPA})_4$ was carried out


under argon and O₂-Ar atmospheres; residues containing varying amts. of BaO, BaO₂ and BaCO₃ were obtained depending on the exptl. conditions.

IT 342805-83-8P

RL: PEP (Physical, engineering or chemical process); PRP (Properties); RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); PROC (Process); RACT (Reactant or reagent)
(polymeric; preparation, crystal structure, sublimation and thermal decomposition)

RN 342805-83-8 HCAPLUS

CN Copper, bis(trimethyl methanetricarboxylato-κO''',κO''')-, (SP-4-1)- (9CI) (CA INDEX NAME)




IT 342805-80-5P

RL: PRP (Properties); RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)
(polymeric; preparation, thermal decomposition and reaction with tri-Me methanetricarboxylate)

RN 342805-80-5 HCAPLUS

CN Copper, methoxy(trimethyl methanetricarboxylato-κO''',κO''')-, (9CI) (CA INDEX NAME)




IT 342805-82-7P

RL: PEP (Physical, engineering or chemical process); PRP (Properties); RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); PROC (Process); RACT (Reactant or reagent)
(preparation, crystal structure, sublimation and thermal decomposition)

RN 342805-82-7 HCAPLUS


CN Copper, bis(tetrahydrofuran)bis(trimethyl methanetricarboxylato-κO''',κO''')-, (OC-6-11)- (9CI) (CA INDEX NAME)



RE.CNT 35 THERE ARE 35 CITED REFERENCES AVAILABLE FOR THIS RECORD
ALL CITATIONS AVAILABLE IN THE RE FORMAT


L4 ANSWER 3 OF 23 HCAPLUS COPYRIGHT 2007 ACS on STN
AN 1998:650912 HCAPLUS
DN 129:312138
TI Copper or zinc complex antimicrobial agents and antimicrobial polymer compositions containing them
IN Kijima, Ichiro; Wakeshima, Ikuko; Ashihara, Wataru; Arikawa, Shinsuke; Ishii, Yoikazu
PA Hoyu System K. K. JP Hoyu Systems K. K., Japan
SO Jpn. Kokai Tokkyo Koho, 8 pp.
CODEN: JKXXAF
DT Patent
LA Japanese
FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	<u>JP 10265310</u>	A	19981006	JP 1997-85506	19970321
PRAI	JP 1997-85506		19970321		
OS	MARPAT 129:312138				
AB	Antimicrobial compns. contain coordination compds. of di- or tricarbonyl compds. $HLC(COR_1)_m(CO_2R_2)_n$ ($1 + m + n = 4$; when $1 = 1$, then the compound is a tricarbonyl compound; when $1 = 2$, then the compound is a dicarbonyl compound; m is preferably ≥ 1) with Cu or Zn ions as active ingredients. The polymer compns. comprise 100 weight parts polymers and 0.001-10 weight parts of the antimicrobial agents. A food container from polypropylene containing 0.05 weight% Zn-acetylacetonate complex (I; preparation given) showed higher antimicrobial effects against Escherichia coli and Staphylococcus aureus than a control containing Mg9ZnO10 instead of I. Applications of the antimicrobial complexes to disinfection of wastewater, fibers, and coatings are also exemplified.				
IT	15378-83-3P, Copper bis(3-ethoxycarbonylacetylacetonate) RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); BUU (Biological use, unclassified); FFD (Food or feed use); SPN (Synthetic preparation); BIOL (Biological study); PREP (Preparation); USES (Uses) (antimicrobial Cu or Zn complexes with carbonyl compds. for polymer compns. and wastewater treatment)				
RN	15378-83-3 HCAPLUS				
CN	Copper, bis[ethyl 2-(acetyl- κ O)-3-(oxo- κ O)butanoato]- (9CI) (CA INDEX NAME)				




- L4 ANSWER 4 OF 23 HCAPLUS COPYRIGHT 2007 ACS on STN
 AN 1998:331599 HCAPLUS
 DN 129:11967
 TI Influence of the Bridging Group of Cross-Conjugated Nitrogenous Bases on the Spectra and Structure of Solvatochromic Mixed-Ligand Copper(II) Chelates Containing β -Ketoenols
 AU Tsiamis, Chris; Hatzidimitriou, Antonis G.; Tzavellas, Leandros C.
 CS Department of Chemistry, University of Thessaloniki, Thessaloniki, 54006, Greece
 SO Inorganic Chemistry (1998), 37(12), 2903-2909
 CODEN: INOCAJ; ISSN: 0020-1669
 PB American Chemical Society
 DT Journal
 LA English
 AB The concomitant reaction of Cu(II) with a heterocyclic nitrogenous cross-conjugated Lewis base (2,2'-dipyridyl ketone, dpk, or 2,2'-dipyridylamine, dpamH) and the anion of a 2-substituted 1,3-dione, β^- , e.g. X-acac $^-$ (X = H, Me, Cl, CN, NO $_2$) affords mixed-ligand chelates. The composition and structure of the chelates depend on the group linking the pyridyl rings. Chelation renders the 2,2'-dipyridyl ketone susceptible to nucleophilic attack by protic mols. It also depends on the group attached to the β -dionato moiety since electron attracting substituents facilitate ligation of a H $_2$ O mol. when the 2,2'-dipyridylamine is present. Spectroscopic observations indicated square pyramidal or distorted tetragonal stereochemistries of the ensuing mixed-ligand chelates with the carbonyl oxygens and the pyridyl nitrogens forming the basal plane. Confirmation was acquired by x-ray structure determination of representative chelates. [Cu(NC-acac)dpamH(H $_2$ O) OClO $_3$] crystallizes in the space group Pnma with Z = 8 (i.e. four mols. per cell), a 17.996(1), b 13.972(1), c 7.801(1) Å. The Cu atom is 2.401(3) Å from the O of the H $_2$ O mol. and 2.60(2) Å from an O atom of the ClO $_4^-$ group. The chelate [Cu(NC-acac)dpC(OH)OCH $_3$ (OClO $_3$)], resulting from the addition of MeOH to the C atom bridging the pyridyl rings, crystallizes in the space group P2 $_1$ /n with a 10.661(1), b 14.987(2), c 13.963(1) Å, β 110.57(3)°, Z = 4. An O of the ClO $_4^-$ group is in apical position, and it is 2.634(8) Å from the Cu atom. The etheric O is distanced 2.615(2) Å from the Cu atom, and the O-Cu-O angle of these weak bonds is only 154.4(2)°. In these chelates the nitrogenous bases adopt the boat conformation with the pyridyl rings forming dihedral angles of .apprx.33°.
 IT 207454-65-7P 207454-72-6P
 RL: PRP (Properties); SPN (Synthetic preparation); PREP (Preparation)
 (preparation and UV-vis. spectrum of)
 RN 207454-65-7 HCAPLUS

CN Copper, aqua(3-nitro-2,4-pentanedionato- $\kappa O, \kappa O'$) (perchlorato- κO) [N-(2-pyridinyl- κN)-2-pyridinamine- $\kappa N1$]-, (OC-6-34)- (9CI) (CA INDEX NAME)



RN 207454-72-6 HCAPLUS

CN Copper, [α -(methoxy- κO)- α -(2-pyridinyl- κN)-2-pyridinemethanol- $\kappa N1$](3-nitro-2,4-pentanedionato- $\kappa O, \kappa O'$)(perchlorato- κO)-, (OC-6-34)- (9CI) (CA INDEX NAME)



RE.CNT 46 THERE ARE 46 CITED REFERENCES AVAILABLE FOR THIS RECORD
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 5 OF 23 HCAPLUS COPYRIGHT 2007 ACS on STN


AN 1998:57648 HCAPLUS

DN 128:148808

TI Axial-equatorial interactions in chelates of β -diones. Crystal and molecular structure of bis(3-nitro-2,4-pentanedionato)copper(II)

AU Tsiamis, Chris; Stergiou, Anagnostis C.; Anesti, Vasileia; Blaton, Norbert

M.; Peeters, Ostwald M.
 CS Department of Chemistry and Department of Physics, University of
 Thessaloniki, Thessaloniki, Greece
 SO Inorganica Chimica Acta (1998), 269(2), 332-336
 CODEN: ICHAA3; ISSN: 0020-1693
 PB Elsevier Science S.A.
 DT Journal
 LA English
 AB The preparation, IR spectrum and structure of bis(3-nitro-2,4-
 pentanedionato)copper(II), [Cu(O₂N-acac)₂], are reported. The structure
 was determined by single crystal x-ray diffraction anal. (triclinic, space
 group P₂1₂1, Z = 1, a 5.841(5), b 7.843(7), c 8.142(8) Å, α
 80.40(7), β 74.59(7), γ 71.80(7)°, R1 = 0.0335, wR2 =
 0.0809). The anion of 3-nitro-2,4-pentanedione, in addition to the commonly
 encountered bonds involving the carbonyl oxygens, also interacts with
 adjacent copper centers through one of the oxygen atoms of the -NO₂ group
 at a distance of 2.609(2) Å and the structure becomes polymeric. The
 axial interactions in the CuO₄2- chromophore are partly responsible for
 the elongation of the in-plane bonds. This is a rare example of a
 bis(β-dionato)copper(II) chelate in which copper(II) exists in a
 rhombically distorted octahedral environment.
 IT 53494-99-8P
 RL: PRP (Properties); SPN (Synthetic preparation); PREP
 (Preparation)
 (preparation, crystal structure and IR spectrum)
 RN 53494-99-8 HCAPLUS
 CN Copper, bis(3-nitro-2,4-pentanedionato-κO,κO')-, (SP-4-1)-
 (9CI) (CA INDEX NAME)



RE.CNT 32 THERE ARE 32 CITED REFERENCES AVAILABLE FOR THIS RECORD
 ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 6 OF 23 HCAPLUS COPYRIGHT 2007 ACS on STN
 AN 1995:875502 HCAPLUS
 DN 123:328439
 TI Hybridization effects on the spectra and structure of solvatochromic
 copper(II) chelates containing β-diones and nitrogenous bases
 AU Tsiamis, Chris; Youri-Tsochatzi, Catherine; Hatzidimitriou, Antony;
 Gourdon, Andre
 CS Department of Chemistry, University of Thessaloniki, Thessaloniki, 54006,
 Greece
 SO Inorganica Chimica Acta (1995), 237(1-2), 93-102
 CODEN: ICHAA3; ISSN: 0020-1693
 PB Elsevier
 DT Journal
 LA English

AB Cu(II) chelates containing a bidentate nitrogenous base (1,2-diamine or heteroarom. α -diimine, enR), the anion of a β -dione, β^- , and the ClO₄⁻ group, [Cu(enR) β]ClO₄, was prepared and characterized and the influence of hybridization of the N donor atoms on the spectra and structure of these compds. was assessed. The spectra (IR, ESR, electronic excitation) indicate that in the absence of steric interference, the CuN2O2 chromophore attains square planar geometry. The ESR observables suggest appreciable covalency. Changes in the structure occur because of the susceptibility of the coordinatively unsatd. [Cu(enR) β]⁺ entity to covalent interactions, which are enhanced as the hybridization of the N donor atoms changes from sp³ to sp². The solvatochromism of the compds. implicates tetragonal distortions in strongly polar solvents and deformed square pyramidal structures in the presence of weakly coordinating anions or mols. Details of the structure of the [Cu(enR) β]⁺ entity and the conformation and configuration of the 1,2-diamines were gathered from the x-ray study of (1-(2-aminoethyl)pyrrolidine)(pentane-2,4-dionato)copper(II) perchlorate, [Cu(aep)acac]ClO₄. The orthorhombic compound crystallizes in the Pna21 space group. The crystal data and the R value are: a 24.5370(7), b 8.4785(4), c 7.6720(3) Å, V_c = 1596(1) Å³, Z = 4, R = 0.0428 for 1099 observed unique reflections.

IT 152748-68-0P 170659-63-9P 170659-65-1P
170659-67-3P 170659-69-5P 170659-71-9P
170659-73-1P

RL: PRP (Properties); SPN (Synthetic preparation); PREP (Preparation)

(preparation and hybridization effects in solvatochromic copper dione nitrogenous base complexes)

RN 152748-68-0 HCAPLUS


CN Copper(1+), (N,N-dimethyl-1,2-ethanediamine-N,N') (3-nitro-2,4-pentanedionato-O2,O4)-, (SP-4-3)-, perchlorate (9CI) (CA INDEX NAME)

CM 1

CRN 152748-67-9

CMF C9 H18 Cu N3 O4


CCI CCS



CM 2

CRN 14797-73-0

CMF Cl O4



RN 170659-63-9 HCAPLUS


CN Copper(1+), (3-nitro-2,4-pentanedionato-O2,O4)(1-pyrrolidineethanamine-N α ,N1)-, (SP-4-3)-, perchlorate (9CI) (CA INDEX NAME)

CM 1

CRN 170659-62-8

CMF C11 H20 Cu N3 O4


CCI CCS



CM 2

CRN 14797-73-0

CMF C1 O4



RN 170659-65-1 HCAPLUS


CN Copper(1+), (1-ethyl-2-pyrrolidinemethanamine-N α ,N1)(3-nitro-2,4-pentanedionato-O2,O4)-, (SP-4-3)-, perchlorate (9CI) (CA INDEX NAME)

CM 1

CRN 170659-64-0

CMF C12 H22 Cu N3 O4


CCI CCS



CM 2

CRN 14797-73-0

CMF Cl O4



RN 170659-67-3 HCAPLUS


CN Copper(1+), (3-nitro-2,4-pentanedionato-O2,O4) (2-pyridinemethanamine-N1,N2)-, (SP-4-3)-, perchlorate (9CI) (CA INDEX NAME)

CM 1

CRN 170659-66-2

CMF Cl1 H14 Cu N3 O4


CCI CCS



CM 2

CRN 14797-73-0


CMF Cl O4



RN 170659-69-5 HCAPLUS
 CN Copper(1+), (2,2'-bipyridine-N,N')(3-nitro-2,4-pentanedionato-O2,O4)-,
 (SP-4-3)-, perchlorate (9CI) (CA INDEX NAME)


CM 1

CRN 170659-68-4
 CMF C15 H14 Cu N3 O4
 CCI CCS



CM 2


CRN 14797-73-0
 CMF C1 O4



RN 170659-71-9 HCAPLUS
 CN Copper(1+), (3-nitro-2,4-pentanedionato-O2,O4)(1,10-phenanthroline-N1,N10)-,
 (SP-4-2)-, perchlorate (9CI) (CA INDEX NAME)

CM 1


CRN 170659-70-8
 CMF C17 H14 Cu N3 O4
 CCI CCS



CM 2

CRN 14797-73-0

CMF Cl O4



RN 170659-73-1 HCAPLUS


CN Copper(1+), (2,9-dimethyl-1,10-phenanthroline-N1,N10) (3-nitro-2,4-pentanedionato-O2,O4)-, perchlorate (9CI) (CA INDEX NAME)

CM 1

CRN 170659-72-0

CMF C19 H18 Cu N3 O4


CCI CCS



CM 2

CRN 14797-73-0

CMF Cl O4



L4 **ANSWER 7 OF 23** HCAPLUS COPYRIGHT 2007 ACS on STN
 AN 1994:123308 HCAPLUS
 DN 120:123308
 TI Counterion effects in the spectra and structure of solvatochromic copper(II) chelates containing 1,2-diamines and β -ketoenols
 AU Tsiamis, Chris; Tzavellas, Leandros C.
 CS Department of Chemistry, University of Thessaloniki, Thessaloniki, 54006, Greece
 SO Inorganica Chimica Acta (1993), 207(2), 179-90
 CODEN: ICHAA3; ISSN: 0020-1693
 DT Journal
 LA English
 AB The concurrent reaction of Cu(II) salts with certain β -diones and N-substituted 1,2-diaminoethanes (enR) is reported. The influence of γ -position derivs. of 2,4-pentanedione (X-acacH; X = H, CN, NO₂) is described, and the contribution of anions to the stability, the stereochem. and the electronic structure of the resulting mixed-ligand chelates is discussed. Spectroscopic observations disclose that the bidentate ligands form chelate rings with Cu(II) as common vertex. [Cu(β -dionato)enR]⁺ are virtually square-planar with CuN₂O₂ chromophore. Cu(II) coordination is unsatd. and in addition to electrostatic interactions that prevail when bulky polyat. anions counterbalance the pos. charge, it is capable of forming covalent bonds with neutral mols. and charged species such as the halides and pseudohalides. The basal CuN₂O₂ plane is distorted upon coordination of unidentate ligands residing

on the apex of the resulting square-pyramidal structure. Further increase in the coordination number of Cu(II) by bidentate anions or neutral mols. leads to tetragonally distorted octahedral structures. This change in the symmetry of the field induced on Cu(II) is revealed by spectral shifts that also disclose covalent interactions in the encounters of the CuN2O2 chromophore with polar or polarizable mols. These interactions are enhanced with increasing ability of the attacking species to act as an electron pair donor. Linear dependence of the ligand field excitation maximum on the donicity of the attacking species was established.

IT 152748-68-0P 152767-34-5P

RL: SPN (Synthetic preparation); PREP (Preparation)
(preparation and solvent effect in electronic spectra of)

RN 152748-68-0 HCAPLUS


CN Copper(1+), (N,N-dimethyl-1,2-ethanediamine-N,N')(3-nitro-2,4-pentanedionato-O2,O4)-, (SP-4-3)-, perchlorate (9CI) (CA INDEX NAME)

CM 1

CRN 152748-67-9

CMF C9 H18 Cu N3 O4


CCI CCS



CM 2

CRN 14797-73-0

CMF Cl O4



RN 152767-34-5 HCAPLUS


CN Copper(1+), (N,N-dimethyl-1,2-ethanediamine-κN,κN')(3-nitro-2,4-pentanedionato-κO,κO')-, (SP-4-3)-, nitrate (9CI) (CA INDEX NAME)

CM 1

CRN 152748-67-9

CMF C9 H18 Cu N3 O4


CCI CCS



CM 2

CRN 14797-55-8

CMF N O3

L4 ANSWER 8 OF 23 HCAPLUS COPYRIGHT 2007 ACS on STN

AN 1993:436490 HCAPLUS

DN 119:36490

TI Substituent and solvent effects in the spectra and structure of some mixed-ligand copper(II) chelates containing β -ketoenols

AU Tsiamis, Chris; Themeli, Maria

CS Department of Chemistry, University of Thessaloniki, Thessaloniki, 54006, Greece

SO Inorganica Chimica Acta (1993), 206(1), 105-15

CODEN: ICHAA3; ISSN: 0020-1693

DT Journal

LA English

AB The IR and the ligand-field excitation spectra of a series of new mixed-ligand copper(II) chelates that encompass N, N-dimethyl-N'-ethyl-ethylenediamine (dmeen) and the anion of a substituted β -ketoenol (1,3-dione) were obtained in the solid state and in solution. Information related to the electronic excitation spectra, the IR spectra, the molar conductivity and the magnetic properties of the newly obtained and characterized chelates are presented and discussed. The molar conductivity in nitromethane reveals a predominance of electrostatic interactions between the $[\text{Cu}(\beta\text{-dione})\text{dmeen}]^+$ entity and bulky polyat. anions that counterbalance the pos. charge while the IR spectra disclose that the bidentate ligands form chelate rings with copper as the common vertex. The resulting CuN_2O_2 chromophore attains a square-coplanar structure and exhibits a tendency for axial ligation which is enhanced when electron-attracting substituents are attached to the β -dionato moiety. The tendency for axial ligation is partially fulfilled when suitable nucleophiles are present. Covalent interactions prevail when chloride is present and upon coordination it presumably occupies the apex of a square-pyramidal structure. Chain-like bidentate anions enable copper(II) to achieve coordination saturation and when configuration requirements demand it, they distort the square-planar arrangement of the

initial CuN2O2 chromophore forming distorted octahedral structures. Covalent solute-solvent interactions are revealed by shifts in the ligand-field excitation spectra that are enhanced as the nucleophilicity of the solvent increases. Linear dependence of the ligand-field excitation maximum on solvent parameters related to donor properties is generally observed

IT 148379-26-4P 148379-27-5P 148379-28-6P

RL: PREP (Preparation)

(preparation and IR spectra and ligand field excitation spectra of)

RN 148379-26-4 HCAPLUS


CN Copper(1+), (N'-ethyl-N,N-dimethyl-1,2-ethanediamine-N,N') (3-nitro-2,4-pentanedionato-O2,O4)-, (SP-4-3)-, perchlorate (9CI) (CA INDEX NAME)

CM 1

CRN 148379-25-3

CMF C11 H22 Cu N3 O4


CCI CCS



CM 2

CRN 14797-73-0

CMF Cl O4



RN 148379-27-5 HCAPLUS


CN Copper(1+), (N'-ethyl-N,N-dimethyl-1,2-ethanediamine-N,N') (3-nitro-2,4-pentanedionato-O2,O4)-, (SP-4-3)-, nitrate (9CI) (CA INDEX NAME)

CM 1

CRN 148379-25-3

CMF C11 H22 Cu N3 O4


CCI CCS



CM 2


CRN 14797-55-8

CMF N O3



RN 148379-28-6 HCAPLUS

CN Copper(1+), (N'-ethyl-N,N-dimethyl-1,2-ethanediamine-N,N')(3-nitro-2,4-pentanedionato-O2,O4)-, chloride, (SP-4-3)- (9CI) (CA INDEX NAME)



● Cl⁻

L4 **ANSWER 9 OF 23** HCAPLUS COPYRIGHT 2007 ACS on STN

AN 1992:50258 HCAPLUS

DN 116:50258

TI Template reactions between bis(3-nitro-2,4-pentanedionato)copper(II) and α,ω-diaminoalkanes in N,N-dimethylformamide solution

AU Fujiwara, Manabu; Wakita, Hisanobu; Matsushita, Takayuki

CS Fac. Sci., Fukuoka Univ., Fukuoka, 814-01, Japan

SO Polyhedron (1991), 10(15), 1773-8

CODEN: PLYHDE; ISSN: 0277-5387

DT Journal

LA English

AB The 1:2 adducts of CuL2 (HL = 3-nitro-2,4-pentanedione) with ethylenediamine, 1,3-diaminopropane, and 1,4-diaminobutane were isolated


and characterized. These adducts are considered to be intermediates in the formation of the corresponding tetraaza macrocyclic complexes and a template reaction mechanism is proposed.

IT 110873-86-4P 138055-62-6P 138072-61-4P

RL: SPN (Synthetic preparation); PREP (Preparation)
(preparation of)


RN 110873-86-4 HCAPLUS

CN Copper, bis(1,2-ethanediamine-N)bis(3-nitro-2,4-pentanedionato-O2,O4)-(9CI) (CA INDEX NAME)




RN 138055-62-6 HCAPLUS

CN Copper, bis(3-nitro-2,4-pentanedionato-O2,O4)bis(1,3-propanediamine-N)-(9CI) (CA INDEX NAME)



RN 138072-61-4 HCAPLUS

CN Copper, bis(1,4-butanediamine-N)bis(3-nitro-2,4-pentanedionato-O2,O4)-(9CI) (CA INDEX NAME)




L4 ANSWER 10 OF 23 HCAPLUS COPYRIGHT 2007 ACS on STN

AN 1990:451421 HCAPLUS


DN 113:51421

TI Preparation and physicochemical studies of some 3-substituted-2,4-pentanedionato copper(II) complexes and their adducts

AU Patel, K. S.; Woods, J. A. O.
 CS Dep. Chem., Univ. Ibadan, Ibadan, Nigeria
 SO Synthesis and Reactivity in Inorganic and Metal-Organic Chemistry (1990),
 20(1), 97-109
 CODEN: SRIMCN; ISSN: 0094-5714
 DT Journal
 LA English
 AB CuL₂ (HL = 3-phenyl-, 3-benzyl-, 3-chloro-, and 3-nitro-2,4-pentanediones)
 and CuL₂.Q (Q = 2,2'-bipyridine (bpy) and 1,10-phenanthroline; n = 1, 2)
 have been prepared and characterized by microanal., conductance, magnetic
 and spectral measurements. The available conductance data in NO₂Me
 indicate that the complexes are nonelectrolytes while the room temperature
 magnetic moments suggest that they are magnetically dilute compds. The
 shifts observed in the ligand field spectral bands and the solvent dependence
 suggest plausible 4-, 5- and 6-coordinate geometries for these compds.
 The IR spectral shifts inferred the possibility of O-unidentate
 β-diketonate linkage and bidentate linkage of base. Square-planar
 and square-pyramidal stereochemistries have been proposed for the
 complexes and adducts, resp., except for Cu(Cl-acac)₂(bpy)₂ (Cl-acacH =
 3-chloroacetylacetone) which is 6-coordinate octahedral.
 IT 14689-25-9P
 RL: RCT (Reactant); SPN (Synthetic preparation); PREP
 (Preparation); RACT (Reactant or reagent)
 (preparation and reaction of, with bipyridine and phenanthroline)
 RN 14689-25-9 HCAPLUS
 CN Copper, bis(3-nitro-2,4-pentanedionato-O₂,O₄)- (9CI) (CA INDEX NAME)



L4 **ANSWER 11 OF 23** HCAPLUS COPYRIGHT 2007 ACS on STN
 AN 1987:589500 HCAPLUS
 DN 107:189500
 TI Template synthesis of a copper(II) complex with tetraaza macrocycle in
 solid phase
 AU Fujiwara, Manabu; Kinoshita, Setsuko; Wakita, Hisanobu; Matsushita,
 Takayuki; Shono, Toshiyuki
 CS Fac. Sci., Fukuoka Univ., Fukuoka, 814-01, Japan
 SO Chemistry Letters (1987), (7), 1323-6
 CODEN: CMLTAG; ISSN: 0366-7022
 DT Journal
 LA English
 GI




AB CuL2(en)2 (HL = 3-nitro-2,4-pentanedione) was prepared in EtOH and characterized by spectroscopic and elemental analyses. The thermal reaction of the adduct in the solid phase resulted in the formation of I by the intramol. condensation.

IT 110873-86-4P, Bis(ethylenediamine)bis(3-nitro-2,4-pentanedionato)copper
 RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)
 (preparation and intramol. cyclocondensation reaction of)

RN 110873-86-4 HCAPLUS

CN Copper, bis(1,2-ethanediamine-N)bis(3-nitro-2,4-pentanedionato-O2,O4)-(9CI) (CA INDEX NAME)



L4 ANSWER 12 OF 23 HCAPLUS COPYRIGHT 2007 ACS on STN

AN 1981:57314 HCAPLUS

DN 94:57314

TI The nucleophilic ring substitution of (3-bromo-2,4-pentanedionato)copper(II)

AU Kasuga, Kuninobu; Nagahara, Takeo; Yamamoto, Yasuo

CS Fac. Sci., Shimane Univ., Matsue, 690, Japan

SO Journal of Coordination Chemistry (1980), 10(4), 217-18
 CODEN: JCCMBQ; ISSN: 0095-8972

DT Journal


LA English

AB Whereas the Br atom of bis(3-bromo-2,4-pentanedionato)copper(II) could not be substituted by nucleophiles such as succinimide, phthalimide, phenol, and benzenethiol, the salts of these nucleophiles produced bis(3-substituted 2,4-pentanedionato)copper(II) complexes.


IT 76401-08-6P
 RL: SPN (Synthetic preparation); PREP (Preparation)
 (preparation of, by nucleophilic substitution reaction of
 bis(3-bromo-2,4-pentanedionato)copper)

RN 76401-08-6 HCAPLUS


CN Copper, bis(3-phenoxy-2,4-pentanedionato-O2,O4)-(9CI) (CA INDEX NAME)




L4 **ANSWER 13 OF 23** HCAPLUS COPYRIGHT 2007 ACS on STN
 AN 1978:114525 HCAPLUS
 DN 88:114525
 TI Nitration of some mixed ligand copper(II) complexes of acetylacetone, benzoylacetone, dibenzoylmethane and Schiff bases
 AU Doraswamy, Uma; Bhattacharya, P. K.
 CS Dep. Chem., Maharaja Sayajirao Univ. Baroda, Baroda, India
 SO Indian Journal of Chemistry, Section A: Inorganic, Physical, Theoretical & Analytical (1977), 15A(9), 828-9
 CODEN: IJCADU; ISSN: 0376-4710
 DT Journal
 LA English
 AB Nitration of the coordinated ligands in complexes of the type $[CuLL']$, where HL = salicylaldimine, 2-hydroxyacetophenonimine, or 2-hydroxy-3-methylacetophenonimide and HL' = acetylacetone, benzoylacetone, or dibenzoylmethane, was carried out. Trinitro compds. were obtained in all cases, except that of 2-hydroxy-3-methylacetophenonimine where a dinitro compound was obtained. The substitution products characterized by elemental anal., magnetic moments, conductance measurements, and IR spectral studies.
 IT 65588-31-0P 65588-32-1P 65888-28-0P
 RL: SPN (Synthetic preparation); PREP (Preparation) (preparation of)
 RN 65588-31-0 HCAPLUS
 CN Copper, [2-(1-iminoethyl)-4,6-dinitrophenolato-N2,O1] (3-nitro-2,4-pentanedionato-O2,O4)- (9CI) (CA INDEX NAME)




RN 65588-32-1 HCAPLUS
 CN Copper, [2-(1-iminoethyl)-4-nitrophenolato-N2,O1] (3-nitro-2,4-pentanedionato-O2,O4)- (9CI) (CA INDEX NAME)




RN 65888-28-0 HCAPLUS
 CN Copper, [2-(iminomethyl)-4,6-dinitrophenolato-N2,O1] (3-nitro-2,4-pentanedionato-O2,O4)- (9CI) (CA INDEX NAME)



L4 ANSWER 14 OF 23 HCAPLUS COPYRIGHT 2007 ACS on STN
 AN 1975:179829 HCAPLUS
 DN 82:179829
 TI Adduct of copper(II) bis(nitroacetylacetonate) with γ -picoline
 AU Mohapatra, B. K.
 CS Dep. Chem., Ravenshaw Coll., Cuttack, India
 SO Chemicke Zvesti (1974), 28(6), 757-9
 CODEN: CHZVAN; ISSN: 0366-6352
 DT Journal
 LA English
 AB Green crystalline $[\text{CuL}_2(\gamma\text{-pic})]$, where HL = nitroacetylacetone and $\gamma\text{-pic}$ = 4-picoline, was prepared and characterized by elec. conductivity, magnetic susceptibility, and ir and electronic spectra.
 IT 55140-20-0P
 RL: SPN (Synthetic preparation); PREP (Preparation)
 (preparation of)
 RN 55140-20-0 HCAPLUS
 CN Copper, (4-methylpyridine)bis(3-nitro-2,4-pentanedionato-O2,O4)- (9CI)
 (CA INDEX NAME)




L4 **ANSWER 15 OF 23** HCAPLUS COPYRIGHT 2007 ACS on STN
 AN 1973:143275 HCAPLUS
 DN 78:143275
 TI Pentacoordinated base adduct of copper(II) nitroacetylacetonate with quinoline
 AU Mohapatra, B. K.
 CS Dep. Chem., Ravenshaw Coll., Cuttack, India
 SO Journal fuer Praktische Chemie (Leipzig) (1973), 315(2), 347-9
 CODEN: JPCEAO; ISSN: 0021-8383
 DT Journal
 LA English
 AB The green crystalline, nonelectrolytic (6.8 mho in Me₂CO), paramagnetic (1 unpaired electron), pentacoordinated 1:1 complex (I) of Cu(II) bis(nitroacetyl-acetonate) with quinoline was prepared and characterized by ir and uv spectra. I was stable in air for a few days. A square pyramidal structure was established for the 1:1 Cu(II) bis(acetyl-acetonate) complex with quinoline (Ooi, S; Fernando, Q., 1967) and a similar structure is expected for I.
 IT 41101-63-7P
 RL: SPN (Synthetic preparation); PREP (Preparation)
 (preparation of)
 RN 41101-63-7 HCAPLUS
 CN Copper, bis(3-nitro-2,4-pentanedionato-O₂,O₄)(quinoline)- (9CI) (CA INDEX NAME)




L4 ANSWER 16 OF 23 HCAPLUS COPYRIGHT 2007 ACS on STN

AN 1972:132200 HCAPLUS 16.923
 DN 76:132200
 TI Mono- and dipyridine adducts of bis(3-substituted 2,4-pentanedionato)copper(II) complexes
 AU Shepherd, T. M.
 CS Dep. Chem., Univ. St. Andrews, St. Andrews, UK
 SO Journal of the Chemical Society, Dalton Transactions: Inorganic Chemistry (1972-1999) (1972), (7), 813-16
 CODEN: JCDTBI; ISSN: 0300-9246
 DT Journal
 LA English
 AB Adduct formation consts., K1, for the equilibrium $\text{CuL}_2 + \text{pyridine} \rightleftharpoons \text{CuL}_2(\text{py})$ (L = AcCRAc; R = Me, H, Cl, Ac, CO₂Et, CN, NO₂) were determined in C₆H₆, Me₂CO, and MeCN at 22°. The K1 values were correlated with the electronic nature of R and the dielec. constant of the solvent. For R = CN and NO₂ further equilibrium $\text{CuL}_2(\text{py}) + \text{pyridine} \rightleftharpoons \text{CuL}_2(\text{py})_2$ were observed in high pyridine concns. and the diadduct formation consts. were determined
 IT 36363-97-0P 36363-99-2P 36364-01-9P
 RL: FORM (Formation, nonpreparative); PREP (Preparation) (formation of)
 RN 36363-97-0 HCAPLUS
 CN Copper, bis(ethyl 2-acetyl-3-oxobutanoato-O₂,O₃)(pyridine)-, (SP-5-21)-(9CI) (CA INDEX NAME)




RN 36363-99-2 HCAPLUS
 CN Copper, bis(3-nitro-2,4-pentanedionato-O₂,O₄)(pyridine)-, (SP-5-21)-(9CI) (CA INDEX NAME)



RN 36364-01-9 HCAPLUS

CN Copper, bis(3-nitro-2,4-pentanedionato-O2,O4)bis(pyridine)- (9CI) (CA INDEX NAME)

L4 **ANSWER 17 OF 23** HCAPLUS COPYRIGHT 2007 ACS on STN

AN 1970:31054 HCAPLUS

DN 72:31054

TI Tautomerism in 3-substituted-2,4-pentanediones and their copper chelates

AU Tanaka, Minoru; Shono, Toshiyuki; Shinra, Koichiro

CS Osaka Univ., Suita, Japan

SO Bulletin of the Chemical Society of Japan (1969), 42(11), 3190-4

CODEN: BCSJA8; ISSN: 0009-2673

DT Journal

LA English


AB Seventeen 3-substituted-2,4-pentanediones were synthesized. The effect of substituents in position 3 on the tautomeric equilibria, i.e., keto-, cis-enol-and trans-enol-forms, has been investigated by NMR spectroscopy. The percentage of each tautomer depends upon the electronic and steric effects of the substituents. The relative intensity of intramol. H bonds in these compds. was evaluated from both the NMR and ir data. The half-wave potential data of polarographic reduction and the Cu-O stretching frequencies of 16 copper chelates of 3-substituted-2,4-pentanediones indicate that the relative stability of the chelate decreases with an increase in the electron-withdrawing effect of the substituents.

IT 15378-83-3P 27970-19-0P


RL: SPN (Synthetic preparation); PREP (Preparation)
(preparation of)

RN 15378-83-3 HCAPLUS


CN Copper, bis[ethyl 2-(acetyl-κO)-3-(oxo-κO)butanoato]- (9CI)
(CA INDEX NAME)




RN 27970-19-0 HCAPLUS
 CN Copper, bis(3-methoxy-2,4-pentanedionato-κO,κO')- (9CI) (CA
 INDEX NAME)



L4 ANSWER 18 OF 23 HCAPLUS COPYRIGHT 2007 ACS on STN
 AN 1968:482976 HCAPLUS
 DN 69:82976
 TI A simple method for preparation of metal acetylacetonates and their nitro
 derivatives
 AU Ebeid, F. M.; Riha, T. I.; Hassanein, M.
 CS Nat. Res. Centre, Dokki, Egypt
 SO Arab Sci. Congr., 5th, Bagdad (1966), Issue Pt. 2, 279-85. Editor(s):
 El-Tahrir, Midan. Publisher: Amer. Univ. at Cairo, Cairo, UAR.
 CODEN: 20ARAH
 DT Conference
 LA English
 AB By mixing the oxides of Al³⁺, Cr³⁺, Cu²⁺, and Mg²⁺ with acetylacetone at
 room temperature in the presence of proper solvents, the corresponding metal
 acetylacetonates were obtained directly in good yield. The solvents used
 for reaction and crystallization were: Et₂O for Al₂O₃; hexane for Cr₂O₃; and
 Me₂CO
 for CuO and MgO. Cr and Cu acetylacetonates undergo nitration with mild
 nitrating agents, e.g. Cu(NO₃)₂·3H₂O. The mol. structures were
 confirmed by elemental anal., ir and uv spectra, m.p., and x-ray
 diffraction methods.
 IT 14689-25-9P
 RL: SPN (Synthetic preparation); PREP (Preparation)
 (preparation of)
 RN 14689-25-9 HCAPLUS
 CN Copper, bis(3-nitro-2,4-pentanedionato-κO,κO')- (9CI) (CA INDEX NAME)




L4 ANSWER 19 OF 23 HCAPLUS COPYRIGHT 2007 ACS on STN
 AN 1967:101194 HCAPLUS
 DN 66:101194
 TI Chemical and spectroscopic studies in metal β -diketonates. II. Nitration of metal β -diketonates.
 AU Singh, Prithwi Raj; Sahai, Ram
 CS Indian Inst. of Technol., Kanpur, India
 SO Australian Journal of Chemistry (1967), 20(4), 649-55
 CODEN: AJCHAS; ISSN: 0004-9425
 DT Journal
 LA English
 AB cf. preceding abstract Bis(3-nitropentane-2,4-dionato)beryllium(II), -copper(II), and -palladium(II); tris(3-nitropentane-2,4-dionato)aluminum(III), -chromium(III), -manganese(III), -iron(III), and -cobalt(III); tetrakis(3-nitropentane-2,4-dionato)zirconium(IV); and tris(3-nitro-1,3-diphenylpropane-1,3-dionato)aluminum(III) and -chromium(III) were prepared by the nitration of the resp. metal chelates of acetylacetone and 1,3-diphenylpropane-1,3-dione. Ferric nitrate, cobaltous nitrate, and nickel nitrate were used as nitrating agents for the 1st time. The nitro chelates were also prepared by a direct, single-step process by using metal nitrates, acetic anhydride, and the ligands. In several cases, metal replacement reactions were observed to be taking place in conjunction with nitration under nitration reaction conditions. Ir, uv, and visible electronic spectra were recorded and are discussed.
 IT 14689-25-9P
 RL: SPN (Synthetic preparation); PREP (Preparation) (preparation of)
 RN 14689-25-9 HCAPLUS
 CN Copper, bis(3-nitro-2,4-pentanedionato-O2,O4)- (9CI) (CA INDEX NAME)



L4 ANSWER 20 OF 23 HCAPLUS COPYRIGHT 2007 ACS on STN
 AN 1967:65003 HCAPLUS
 DN 66:65003

TI Simple method for preparation of metal acetylacetonates and their nitro derivatives 20 223
 AU Ebeid, Fathi M.; Rihan, T. I.; Hassanein, M.
 CS Natl. Res. Centre, Cairo, Egypt
 SO Indian Journal of Chemistry (1966), 4(10), 451-2
 CODEN: IJOCAP; ISSN: 0019-5103
 DT Journal
 LA English
 GI For diagram(s), see printed CA Issue.
 AB Metal acetylacetonates (Ia) were obtained directly in good yields by mixing MeCQCH₂COMe with an excess of a metal oxide at room temperature in the presence of a proper redistd. solvent. The rate of reaction could be accelerated by refluxing. Following are given the metal oxide used, reaction medium, and m.p. of the corresponding Ia: Al₂O₃, ether, 193° (ether) (λ 280 mμ); Cr₂O₃, hexane, 214-15° (hexane) (I) (λ 255-272, 335 mμ); CuO, Me₂CO, -(sublimed) (II) (λ 293 mμ); MgO, Me₂CO, 266-7° (λ 280 mμ). I and II were nitrated by using Cu(NO₃)₂·3H₂O in Ac₂O to yield the corresponding nitro derivs., m. 245° (ν 1200 cm.⁻¹), and 193° (ν 1675 cm.⁻¹), resp. Usual nitrating agents destroyed Ia. The results were in accordance with those of Collmann, et al. (CA 55, 9274c) and consequently confirmed the aromatic character of Ia.
 IT 14689-25-9P
 RL: SPN (Synthetic preparation); PREP (Preparation) (preparation of)
 RN 14689-25-9 HCAPLUS
 CN Copper, bis(3-nitro-2,4-pentanedionato-O₂,O₄)- (9CI) (CA INDEX NAME)



L4 ANSWER 21 OF 23 HCAPLUS COPYRIGHT 2007 ACS on STN
 AN 1965:454744 HCAPLUS
 DN 63:54744
 OREF 63:9975d-g
 TI Aluminum nitrate and nitric acid as nitrating agents for acetyl-acetonates
 AU Sen, D. N.; Thankarajan, N.
 CS Nat. Chem. Lab., Poona
 SO Indian Journal of Chemistry (1965), 3(5), 215-16
 CODEN: IJOCAP; ISSN: 0019-5103
 DT Journal
 LA English
 AB Tris(3-nitro-2,4-pentanediono)aluminum(I) was prepared in a single step process, by using Al(NO₃)₃ and acetylacetone (Ia) in Ac₂O. Al(NO₂)₂ was also employed to prepare tris(3-nitro-2,4-pentane-diono)beryllium(II), which could not be prepared with Cu(NO₃)₂. Concentrated HNO₃ was a good nitrating agent for the metals acetyl-acetonates. Thus, 3.75 g. powdered Al(NO₃)₃·9H₂O was

added (30 min.) in 6 portions to a stirred and cooled mixture of 3 ml. Ia in 20 ml. Ac2O. The mixture was stirred 4 hrs. at room temperature, care being taken

to see that the temperature did not rise. The solution was poured into .apprx.200


ml. ice-cold H2O containing NaOAc and left 1 hr. to give 2.3 g. I, m. 250° (C6H66). Powdered Al(NO3)3 (3.75 g.) was added slowly (as for I) to 3.5 g. Be acetylacetamato and 25 ml. Ac2O. Working up the reaction mixture yielded 2 g. II, m. 185.5°. Bis(3-nitro-2,4-pentanediono)copper (III) and tris(3-nitro-2,4-pentanediono)chromium(IV), m. 255°, were prepared in the same way as above from the resp. metal acetylacetonate, Al(NO3)3 and Ac2O. Attempted nitration of Be p-acetylacetonate using Cu(NO3)2 led to the isolation of III. Powdered Be acetylacetonate (3.1 g.) was added slowly (1 hr.) to a cooled and stirred solution of 3.7 g. Cu(NO3)2.3H2O in 30 ml. Ac2O. The solution was stirred 2 hrs. at room temperature and poured into cold-H2O to yield 3 g. III, m. 230-4° (decomposition), (40:60 EtOH-CHCl3). Concentrated HNO3 (16N; 2.2 ml.) was added dropwise to a cooled (0°) and stirred suspension of 3.25 g. Al acetylacetonate in 25 ml. Ac2O. The solution was stirred 1 hr. more at 0° and then 4 hrs. at room temperature. Working up yielded 2.4 g. I. A discussion of nitrating agents and conditions is presented.

IT 14689-25-9P, Copper, bis(3-nitro-2,4-pentanedionato)-

RL: PREP (Preparation)
(preparation of)

RN 14689-25-9 HCAPLUS

CN Copper, bis(3-nitro-2,4-pentanedionato-O2,O4)- (9CI) (CA INDEX NAME)



L4 ANSWER 22 OF 23 HCAPLUS COPYRIGHT 2007 ACS on STN

AN: 1964:7612 HCAPLUS

DN 60:7612

OREF 60:1311f

TI Preparation and reduction of bis-(3-nitro-2,4-pentanedione)-beryllium

AU Klein, Richard M.; Bailar, John C., Jr.

CS Univ. of Illinois, Urbana

SO Inorg. Chem. (1963), 2(6), 1187-90


DT Journal

LA Unavailable

AB The reaction of 2,4-pentanedione with Be nitrate trihydrate in Ac2O gave bis-(3-nitro-2,4-pentanediono)beryllium. This was reduced catalytically to the corresponding amine. Bis-(2,4-pentanediono)beryllium reacted in this nitrating medium to yield a polymeric nitrated complex containing hydroxo bridges. This was converted to either the methoxo or ethoxo bridged complexes by boiling in MeOH or EtOH. These were shown by mol. weight studies to be partially trimeric.

IT 14689-25-9P, Copper, bis(3-nitro-2,4-pentanedionato)-

RL: PREP (Preparation)
 (preparation of)
 RN 14689-25-9 HCAPLUS
 CN Copper, bis(3-nitro-2,4-pentanedionato-O2,O4)- (9CI) (CA INDEX NAME)



L4 **ANSWER 23 OF 23** HCAPLUS COPYRIGHT 2007 ACS on STN

AN 1963:18976 HCAPLUS

DN 58:18976

OREF 58:3090h,3091a-b

TI The reactivity of coordinated acetylacetone

AU Djordjevic, C.; Lewis, J.; Nyholm, R. S.

CS Univ. Coll., London

SO Journal of the Chemical Society (1962) 4778-84

CODEN: JCSOA9; ISSN: 0368-1769

DT Journal

LA Unavailable

AB The reactivity of acetylacetone, coordinated to a metal atom, with various reagents was investigated. The complexes of Cu, Ni, Pt, and Al yield γ -nitro-substituted complexes with N2O4. NOCl yields nitroso or chloro complexes, depending on the conditions employed. Nitrite ions in the presence of NH3 at pH 7 yield iminoacetylacetone derivs. of Ni and Pd; these are typical diamagnetic, square complexes. IR spectra were used extensively to identify the substituted groups present and to assign structures to the complexes isolated. Several unusual types of synthetic organic reactions dependent on the fact that the acetylacetone is coordinated to a metal atom were identified.


IT 14689-25-9P, Copper, bis(3-nitro-2,4-pentanedionato)-

RL: PREP (Preparation)

(preparation of)

RN 14689-25-9 HCAPLUS

CN Copper, bis(3-nitro-2,4-pentanedionato-O2,O4)- (9CI) (CA INDEX NAME)



10529569 FlourineFreeMetallicComplexes for GasPhaseChemDepo

=> dis his

(FILE 'HOME' ENTERED AT 17:51:45 ON 16 JAN 2007)

FILE 'REGISTRY' ENTERED AT 17:51:56 ON 16 JAN 2007

L1 STRUCTURE UPLOADED

L2 2 S L1 SAM

L3 49 S L1 FULL

FILE 'HCAPLUS' ENTERED AT 17:55:47 ON 16 JAN 2007

L4 23 S L3/PREP

=> save l1-l4

ENTER NAME OR (END):fluorstr/l

L# LIST L1-L4 HAS BEEN SAVED AS 'FLUORSTR/L'